

## Estimation of CO<sub>2</sub> partial pressure and bicarbonate concentration in the hemolymph of the noble scallop *Mimachlamys nobilis*

メタデータ	言語: English 出版者: 水産大学校 公開日: 2024-10-11 キーワード (Ja): キーワード (En): <i>Mimachlamys nobilis</i> ; cannulation; hemolymph; acid-base balance; Pco <sub>2</sub> ; dissociation constant of carbonic acid 作成者: 半田, 岳志, 山元, 憲一 メールアドレス: 所属: 水産研究・教育機構
URL	<a href="https://fra.repo.nii.ac.jp/records/2012072">https://fra.repo.nii.ac.jp/records/2012072</a>

This work is licensed under a Creative Commons Attribution 4.0 International License.



## Estimation of CO<sub>2</sub> partial pressure and bicarbonate concentration in the hemolymph of the noble scallop *Mimachlamys nobilis*

Takeshi Handa<sup>†</sup> and Ken-ichi Yamamoto

**Abstract :** We examined hemolymph pH, total CO<sub>2</sub> content (Tco<sub>2</sub>, mM/l), CO<sub>2</sub> partial pressure (Pco<sub>2</sub>, mmHg) and bicarbonate concentration ([HCO<sub>3</sub><sup>-</sup>], mM/l) in order to evaluate the acid-base balance of the noble scallop *Mimachlamys nobilis* in normoxic conditions. Hemolymph was collected anaerobically through a cannula after catheterization of the adductor muscle. The mean values of hemolymph pH and Tco<sub>2</sub> were 7.442 and 1.50 mM/l, respectively. Using the CO<sub>2</sub> solubility coefficient and apparent dissociation constant of carbonic acid determined in this study, Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] were calculated as 1.55 mmHg and 1.44 mM/l, respectively. These values were in same range as those of the hemolymph of other marine bivalves of Mytiloidea, Pterioidea and Ostreoida.

**Key words :** *Mimachlamys nobilis*, cannulation, hemolymph, acid-base balance, Pco<sub>2</sub>, dissociation constant of carbonic acid

### Introduction

The noble scallop *Mimachlamys nobilis* is a filibranchial bivalve classified in the Pectinidae, Ostreoida.<sup>1)</sup> This species inhabits rock reefs or sandy-bottomed shores up to a depth of 20 m in the littoral zone from the Boso Peninsula to Okinawa in Japan along the Kuroshio and Tsushima currents.<sup>1-2)</sup> These warm ocean currents maintain a moderate climate across this region, with annual water temperatures of 17–29°C.<sup>2)</sup> Noble scallops are produced in southwest Japan, including Wakayama, Ohita, Nagasaki, Kagoshima and Okinawa prefectures, and are importance species for aquaculture.<sup>1-3)</sup> The many species of the Pectinidae distributed around the world are widely found in high latitudes between 30° and 55° in the Northern and Southern hemispheres and are economically important aquatic resources in cold-water sea areas.<sup>4)</sup> The noble scallop is, therefore, an unusual resource as a Pectinidae species produced in warm-water sea areas. Noble scallop has been a subject of previous research in terms of reproductive cycle,<sup>5)</sup> induction of oviposition,<sup>6)</sup> seedling production,<sup>6)</sup> early food,<sup>7)</sup> karyotype,<sup>8)</sup> genetic variation,<sup>9)</sup> gametogenesis and triploid induction.<sup>10,11)</sup> The

ciliary movement of the ctenidium in hypoxic and anathermal conditions has been studied.<sup>12)</sup> The anatomical structures of the ctenidium and circulatory system were clarified recently.<sup>13)</sup> However, there are few reports on the respiratory mechanism from the viewpoint of CO<sub>2</sub> dynamic phase and acid-base balance. Research into the acid-base status could contribute to efficient CO<sub>2</sub> utilization, which is related to respiration, and calcification for the formation of the shell valves. The estimated CO<sub>2</sub> partial pressure of the hemolymph was 0.9 mmHg in sea mussel *Mytilus edulis*,<sup>14)</sup> 2.3 mmHg in Asian freshwater clam *Corbicula fluminea*,<sup>15)</sup> 1.7–2.3 mmHg in akoya pearl oyster *Pinctada fucata*,<sup>16,17)</sup> and 1.5 mmHg in black-lip pearl oyster *Pinctada margaritifera*.<sup>18)</sup> Because the CO<sub>2</sub> partial pressures in these bivalves were very low, it was supposed that the CO<sub>2</sub> partial pressure in the noble scallop would also be similarly very low. The direct measurement of CO<sub>2</sub> partial pressure is difficult when there is small quantity of hemolymph sample and CO<sub>2</sub>. The estimation of CO<sub>2</sub> partial pressure by application of the Henderson–Hasselbalch equation is used in studies of the acid-base balance owing to the relative ease and accuracy of the estimates.<sup>19)</sup> In the equation, the characteristic values of the CO<sub>2</sub>

solubility coefficient ( $\alpha\text{CO}_2$ ) and apparent dissociation constant of carbonic acid (pKapp) in the hemolymph are required for the experimental animal. Therefore, we examined noble scallop hemolymph pH, total  $\text{CO}_2$  content,  $\text{CO}_2$  partial pressure and bicarbonate concentration using the hemolymph  $\alpha\text{CO}_2$  and pKapp, which were determined in this study. By means of catheterization of the adductor muscle, hemolymph was collected anaerobically from noble scallop underwater.

## Materials and Methods

### Experimental animals and conditions

The experiments used 36 noble scallops (shell length:  $86.1 \pm 5.0$  mm [Mean  $\pm$  SE]), shell height:  $88.0 \pm 5.1$  mm, and total wet weight:  $102.7 \pm 11.8$  g). The animals were obtained from the coast of Nagasaki prefecture, Japan. After cleaning the shell valves, they were reared for 2 months at  $24^\circ\text{C}$  in aerated seawater with added cultivated phytoplankton.<sup>17,18,20,21</sup> Twenty-four hours before collecting hemolymph, the noble scallop were transferred to seawater that was particle-free ( $>0.45 \mu\text{m}$ ). All experiments were conducted in the seawater with a salinity of 33 psu, water temperature  $24.0^\circ\text{C}$ ,  $\text{O}_2$  saturation 99%, pH 8.15, and total  $\text{CO}_2$  content 1.2 mM/l.

### Surgical procedures and hemolymph collection

Hemolymph was collected from the adductor muscle using a cannula (polyethylene tubing, 0.96 mm outer diameter, 0.58 mm inner diameter, PE-50, Clay Adams). A small hole (2 mm diameter) was made on adjacent shell valves at the center of the posterior margin. The cannula with a stylet was inserted through the hole into the adductor muscle, which is part of the large anterior muscle, and was advanced 1.0 – 1.2 cm toward the center of the adductor muscle. The stylet was removed, and the end of the cannula was closed. The cannula was gently fixed to the left shell valve using denture adhesive (Kobayashi Pharmaceutical Co., Ltd.) in order to prevent effects from movement of the shell valves. This surgical operation was completed within 5 minutes. The cannulated scallop was transferred to a darkened acrylic respiratory chamber and was allowed to recover for 3 h at  $24.0^\circ\text{C}$  in normoxic conditions. A hemolymph sample was then drawn through the cannula using a gas-tight micro syringe (Model 1750, Hamilton Co. Ltd.). The

volume of hemolymph collected was 0.4 ml.

### Hemolymph analysis

Hemolymph pH and total  $\text{CO}_2$  content ( $\text{Tco}_2$ , mM/l) were measured immediately after each collection. The pH was measured using a blood gas meter (BGM200; Cameron Instruments), pH glass and reference electrodes (E301, E351; Cameron Instruments) at  $24.0^\circ\text{C}$ .  $\text{Tco}_2$  was measured using a total  $\text{CO}_2$  analyzer (Capnicon 5; Cameron Instruments). The hemolymph  $\text{CO}_2$  partial pressure ( $\text{Pco}_2$ , mmHg) and bicarbonate concentration ( $[\text{HCO}_3^-]$ , mM/l) were calculated by rearranging the Henderson–Hasselbalch equation.<sup>19,22</sup> In the equation, the  $\text{CO}_2$  solubility coefficient ( $\alpha\text{CO}_2$ ,  $\mu\text{M}/\text{mmHg}$ ) and apparent dissociation constant of carbonic acid (pKapp) of noble scallop were required. The determinations of  $\alpha\text{CO}_2$  and pKapp were performed *in vitro*.

The  $\alpha\text{CO}_2$  was determined using noble scallop hemolymph that was adjusted to pH 2.5 by the addition of lactic acid (Wako Pure Chemical Industries, Ltd.). The acidified sample was transferred to a tonometer flask and equilibrated using humidified standard  $\text{CO}_2$  gas ( $\text{CO}_2$ , 15.0%;  $\text{O}_2$ , 20.9%;  $\text{N}_2$  Balance) using an equilibrator (DEQ-1; Cameron Instruments) at  $24.0^\circ\text{C}$ , and subsequently the total  $\text{CO}_2$  content of each equilibrated sample was measured with the total  $\text{CO}_2$  analyzer. The  $\text{CO}_2$  partial pressure of the equilibrated sample was calculated from known  $\text{CO}_2$  concentration standard gas (15.0%), prevailing barometric pressure and water vapor pressure at  $24.0^\circ\text{C}$ . The  $\text{CO}_2$  solubility coefficient was calculated using the equation:

$$\alpha\text{CO}_2 = \text{Total CO}_2 \text{ content} \cdot \text{CO}_2 \text{ Partial pressure}^{-1}$$

For determination of the apparent dissociation constant of carbonic acid, hemolymph was transferred to a tonometer flask and equilibrated with humidified standard  $\text{CO}_2$  gases ( $\text{CO}_2$ , 0.5, 1.0, 2.0, 5.0 and 10.0%;  $\text{O}_2$ , 20.9%;  $\text{N}_2$  Balance) using an equilibrator at  $24.0^\circ\text{C}$ . After equilibration, the pH and total  $\text{CO}_2$  content of the sample were measured with the blood gas meter and the total  $\text{CO}_2$  analyzer. Using the sample pH, total  $\text{CO}_2$  content and  $\alpha\text{CO}_2$  calculated using the above equation, the pKapp was determined by rearrangement of Henderson–Hasselbalch equation<sup>19,22</sup> as follows:

$$pK_{app} = pH - \log \left[ \frac{\text{Total CO}_2 \text{ concentration} - \alpha\text{CO}_2 \cdot \text{CO}_2 \text{ Partial pressure}}{\alpha\text{CO}_2 \cdot \text{CO}_2 \text{ Partial pressure}} \right]^{-1}$$

where CO<sub>2</sub> partial pressure is calculated from known CO<sub>2</sub> concentration standard gases.

The  $\alpha\text{CO}_2$  and pK<sub>app</sub> obtained in this study were used for calculation of hemolymph Pco<sub>2</sub> from measured pH and Tco<sub>2</sub>:

$$P_{\text{CO}_2} = T_{\text{CO}_2} \cdot [\alpha\text{CO}_2 \cdot (1 + 10^{(pH - pK_{app})})]^{-1}$$

[HCO<sub>3</sub><sup>-</sup>] was calculated from Tco<sub>2</sub>,  $\alpha\text{CO}_2$  and Pco<sub>2</sub> using the following equation<sup>23</sup>:

$$[\text{HCO}_3^-] = T_{\text{CO}_2} - \alpha\text{CO}_2 \cdot P_{\text{CO}_2}$$

#### Statistical analysis

All data are expressed as means  $\pm$  standard error. Kruskal-Wallis one-way analysis of variance was used to test for changes in hemolymph properties using the standard CO<sub>2</sub> gases. Mann-Whitney *U* test was used for the comparison of mean values of hemolymph parameters. Statistically significant differences were set at  $P < 0.01$ .

## Results

Hemolymph samples were collected from the adductor muscles of noble scallops through cannulae. The collection volume was 0.4–0.5 ml from each individual. The hemolymph pH and Tco<sub>2</sub> in normoxic conditions were  $7.442 \pm 0.0094$  and  $1.50 \pm 0.043$  mM/l, respectively (Table 1), and the hemolymph  $\alpha\text{CO}_2$  was  $39.0 \pm 0.820$   $\mu\text{M}/\text{mmHg}$ . The hemolymph pK<sub>app</sub> at known CO<sub>2</sub> partial pressures (standard gases) and the corresponding measured pH and Tco<sub>2</sub> values are shown in Table 2. The calculated pK<sub>app</sub> from all hemolymph samples was  $6.064140 \pm 0.013876$ . Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] were calculated by substitution of the mean value of hemolymph  $\alpha\text{CO}_2$  and pK<sub>app</sub> in the rearranged Henderson–Hasselbalch equation as follows:

$$P_{\text{CO}_2} = T_{\text{CO}_2} \cdot [0.039 \cdot (1 + 10^{(pH - 6.064140)})]^{-1}$$

$$[\text{HCO}_3^-] = T_{\text{CO}_2} - 0.039 \cdot P_{\text{CO}_2}$$

where the units of the parameters in the equations are mmHg for Pco<sub>2</sub>, mM/l for Tco<sub>2</sub> and for [HCO<sub>3</sub><sup>-</sup>].

Hemolymph Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] at 24.0°C in normoxic condition were  $1.55 \pm 0.070$  mmHg and  $1.44 \pm 0.041$  mM/l, respectively

**Table 1.** Hemolymph pH, total CO<sub>2</sub> content (Tco<sub>2</sub>), CO<sub>2</sub> partial pressure (Pco<sub>2</sub>) and bicarbonate concentration ([HCO<sub>3</sub><sup>-</sup>]) of noble scallop (*Mimachlamys nobilis*) at 24°C in normoxic conditions

	Mean	SE	N
pH	7.442	0.0094	6
Tco <sub>2</sub> mM/l	1.50	0.043	6
Pco <sub>2</sub> mmHg	1.55	0.070	6
[HCO <sub>3</sub> <sup>-</sup> ] mM/l	1.44	0.041	6

$\alpha\text{CO}_2$ , 39.01  $\mu\text{M}/\text{mmHg}$ ; pK<sub>app</sub>, 6.0641396  $\pm$  0.0138764


**Table 2.** Mean values of measured pH, total CO<sub>2</sub> content (Tco<sub>2</sub>) and calculated apparent dissociation constant of carbonic acid (pK<sub>app</sub>) of hemolymph in the adductor muscle of the noble scallop (*Mimachlamys nobilis*) with known Pco<sub>2</sub> standard gases

Standard gas		Hemolymph			
CO <sub>2</sub> (%)	Pco <sub>2</sub> (mmHg)	pH	Tco <sub>2</sub> (mM/l)	pK <sub>app</sub>	N
0.5	3.64	7.213	1.791	6.1532	6
1.0	7.28	6.873	1.937	6.1089	6
2.0	14.60	6.629	2.766	6.0495	6
5.0	36.40	6.265	4.104	6.0111	6
10.0	72.80	5.990	5.973	5.9980	6

Water temperature 24.0°C; Barometric pressure 759 mmHg; Water vapor pressure 22.37 mmHg

(Table 1). In in vitro experiments (Table 2), the pH decreased significantly with the increase in  $P_{\text{CO}_2}$  ( $P < 0.01$ ). At the same time, the interaction between pKapp and pH was analyzed (Fig. 1), and the correction equation of pKapp was obtained as follows:

$$\text{pKapp} = 45.217 - 17.721 \cdot \text{pH} + 2.6463 \cdot \text{pH}^2 - 0.1303 \cdot \text{pH}^3$$



**Fig 1.** Relationship between pH and apparent dissociation constant of carbonic acid (pKapp) in the hemolymph of noble scallop *Mimachlamys nobilis* at 24°C (n=30). Values are means  $\pm$  SE. The solid line is fitted to the data and the equation:  $\text{pKapp} = 45.217 - 17.721 \cdot \text{pH} + 2.6463 \cdot \text{pH}^2 - 0.1303 \cdot \text{pH}^3$  ( $R^2=0.992$ )

For the comparison, using the hemolymph pH and  $T_{\text{CO}_2}$  measured immediately after collection,  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  were estimated using the correction equation and are shown in Table 3. The hemolymph  $P_{\text{CO}_2}$  calculated using the correction equation was higher than that from the mean value of pKapp ( $P < 0.01$ ). There was no difference in hemolymph  $[\text{HCO}_3^-]$  calculated by the two methods (Table 3).

## Discussion

We collected noble scallop hemolymph from the adductor muscle and examined hemolymph pH, total  $\text{CO}_2$  content,  $\text{CO}_2$  partial pressure and bicarbonate concentration in order to evaluate the acid-base balance of noble scallops. The hemolymph was collected anaerobically through a cannula from noble scallops kept underwater after pretreatment by adductor muscle catheterization. The mean values of pH and  $T_{\text{CO}_2}$  measured immediately after hemolymph collection were 7.442 and 1.50 mM/l, respectively. Previously reported mean values of hemolymph pH include 7.65 in sea mussel *M. edulis* at 12°C,<sup>14)</sup> 7.36 in Pacific oyster *Crassostrea gigas* at 15°C,<sup>24)</sup> 7.55 in *M. galloprovincialis* at 18°C,<sup>25)</sup> 7.284–7.375 in akoya pearl oyster *P. fucata* at 28°C,<sup>17)</sup> and 7.563 in black-lip pearl oyster *P. margaritifera* at 26°C.<sup>18)</sup> Although there are few descriptions of hemolymph total  $\text{CO}_2$  content in marine bivalves, Handa and Yamamoto (2011, 2012, 2015) reported the mean values in akoya pearl oyster *P. fucata* and black-lip pearl oyster *P. margaritifera* as 1.90–2.10 mM/l<sup>16,17)</sup> and 2.04 mM/l,<sup>18)</sup> respectively. The hemolymph pH in noble scallop was almost the same as that in other marine bivalves, and the contents of carbonic acid and  $\text{CO}_2$  in noble scallop hemolymph appeared to be less than in pearl oysters.

Cameron (1986) reported the  $\text{CO}_2$  solubility of seawater as a function of temperature, and the solubility coefficient at 24°C was 39.24–40.13  $\mu\text{M}/\text{l}/\text{mmHg}$  at 30–35 salinity (psu).<sup>26)</sup> The hemolymph  $\alpha\text{CO}_2$  in noble scallop was in the range of the coefficient of seawater. The mean value of hemolymph pKapp in this study was 6.0641396, whereas the hemolymph pKapp

**Table 3.** Comparison of the hemolymph  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  calculated from the mean value of pKapp and using the correction equation

	$P_{\text{CO}_2}$ (mmHg)	$[\text{HCO}_3^-]$ (mM/l)	N
the mean value of pKapp	1.55	1.44	6
the correction equation of the pKapp	2.05 *	1.42	6

\* : statistically significant difference (Mann-Whitney *U* test,  $P < 0.01$ )

values of other marine bivalves were 5.8191 in the akoya pearl oyster *P. fucata* at 28°C,<sup>17)</sup> 5.99878 in black-lip pearl oyster *P. margaritifera* at 26°C,<sup>18)</sup> and 6.114 in sea mussel *M. edulis* at 12°C.<sup>14,27)</sup> The apparent dissociation constant of carbonic acid is equal to the pH at which it shows the most effective as a buffer.<sup>28)</sup> The most effective buffer pH of noble scallop hemolymph seemed to be lower than that in sea mussel *M. edulis* but higher than that in pearl oyster *P. fucata*.

Using the hemolymph  $\alpha\text{CO}_2$  and pKapp in this study, Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] of the hemolymph of noble scallops were calculated. The mean values of hemolymph Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] were 1.55 mmHg and 1.44 mM/l, respectively (Table 1). In other marine bivalves, the mean values of hemolymph Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] were 0.9 mmHg and 1.8 mM/l in sea mussel *M. edulis* at 12°C,<sup>14)</sup> 0.15 kPa (2.0 mmHg) and 1.37 mM/l in Pacific oyster *C. gigas* at 15°C,<sup>24)</sup> 1.15 mmHg and 1.62 mM/l in *M. galloprovincialis* at 18°C,<sup>25)</sup> 2.08–2.33 mmHg and 1.83–2.04 mM/l in the akoya pearl oyster *P. fucata* at 28°C,<sup>17)</sup> and 1.50 mmHg and 1.98 mM/l in black-lip pearl oyster *P. margaritifera*,<sup>18)</sup> respectively. The acid-base status of the noble scallop will be almost same as it of those bivalves regardless of water temperature.

In decapod crustaceans, the hemolymph Pco<sub>2</sub> was 0.21 kPa (1.57 mmHg) in green crab *Carcinus maenas*, 0.32 kPa (2.4 mmHg) in blue crab *C. sapidus*, 0.28–0.46 kPa (2.1–3.45 mmHg) in Chinese mitten crab *Eriocheir sinensis*,<sup>29,30)</sup> 0.44 kPa (3.3 mmHg) in European lobster *Homarus gammarus*,<sup>31)</sup> and 0.268 kPa (2.01 mmHg) in American lobster *H. americanus*.<sup>32)</sup> The hemolymph [HCO<sub>3</sub><sup>-</sup>] of these decapods were 5.5 mM/l in green crab *Carcinus maenas*, 4.2 mM/l in blue crab *C. sapidus*, 7.09–8.0 mM/l in Chinese mitten crab *Eriocheir sinensis*,<sup>29,30)</sup> and 9.2–9.3 mM/l in European lobster *H. gammarus*.<sup>31)</sup> In teleosts, blood Pco<sub>2</sub> was 2.0 mmHg in red sea bream *Pagrus major*,<sup>33)</sup> 2.0–4.0 mmHg in starry flounder *Platichthys stellatus*,<sup>34)</sup> 0.42–0.48 kPa (3.2–3.6 mmHg) in Atlantic cod *Gadus morhua*,<sup>35)</sup> 3.47–4.98 mmHg in rainbow trout *Salmo gairdneri* (*Oncorhynchus mykiss*),<sup>34)</sup> and 3.8–4.2 mmHg in the common carp *Cyprinus carpio*.<sup>36,37)</sup> The [HCO<sub>3</sub><sup>-</sup>] was 8.3–9.2 mM/l in Atlantic cod *G. morhua*,<sup>35)</sup> 8.8–9.3 mM/l in rainbow trout *S. gairdneri* (*O. mykiss*),<sup>34)</sup> and 10.2–10.3 mM/l in the common carp *C. carpio*.<sup>37)</sup> Marine bivalves could contain CO<sub>2</sub> and bicarbonate levels that are less than crustacean and teleosts; therefore, noble scallop seemed to utilize CO<sub>2</sub> for calcification of shell valves and discharge CO<sub>2</sub> to control the acid-base balance

of the hemolymph.

On the other hand, the pH and Tco<sub>2</sub> changed significantly with the increase in Pco<sub>2</sub> (Table 2). The relationship of hemolymph pH and pKapp was shown (Fig. 1), and Pco<sub>2</sub> and [HCO<sub>3</sub><sup>-</sup>] were estimated on the basis of the correction equation. Hemolymph Pco<sub>2</sub> estimated by the correction equation was higher than that from the mean value of pKapp, nevertheless there was no significant difference in [HCO<sub>3</sub><sup>-</sup>] (Table 3). The  $\alpha\text{CO}_2$  and pKapp vary with ionic strength and temperature,<sup>19)</sup> and estimation of Pco<sub>2</sub> could be affected by temperature and salinity. The hemolymph pH of noble scallops in normoxic conditions in this study (7.442, Table 1) was higher than the *in vitro* pH range (5.990–7.213, Table 2). Therefore, it is necessary to examine these parameters in various temperature, salinity and pH conditions in order to increase the accuracy of the calculation of Pco<sub>2</sub> and to formulate the correction equation for pH, salinity and water temperature.

## References

- Okutani T: Pterioidea. In: Okutani T (ed) Marine Mollusks in Japan. Tokai University Press, Tokyo, 902-903 (2000)
- Nogami K: Recent developments in shellfish culture in southern Japan. *NOAA Technical report NMFS*, **16**, 73-75 (1984)
- Ito H: *Chlamys (Mimachlamys) senatrix nobilis*. In: Shumway SE (ed) Scallops: Biology, Ecology and Aquaculture, Developments in Aquaculture and Fisheries Science, **21**. Elsevier Science Publishers, 1051 (1991)
- Brand RA: Scallop ecology, distributions and behaviour. In: Shumway SE (ed) Scallops: Biology, Ecology and Aquaculture, Developments in Aquaculture and Fisheries Science, **21**. Elsevier Science Publishers, 517-584 (1991)
- Komaru A and Wada KT: Seasonal changes of gonad cultured scallops, *Chlamys nobilis*. *Bull Nat Res Inst Aquaculture*, **14**, 125-132 (1988)
- Hirata H, Shinomiya A, Kadowaki S, Nakazono T and Kasedo T: Seed production of scallop *Chlamys nobilis* – I, Spawning induced by dark stocking. *Mem Fac Fish Kagoshima Univ*, **27**, 289-294 (1978)
- Kadowaki S, Nakazono T, Ioku S, Kasedo T and Hirata H: Seed production of scallop *Chlamys nobilis* – II, Mixture diet

- of marine yeast and *Chlorella* sp. for the veliger larvae. *Mem Fac Fish Kagoshima Univ*, **29**, 209-215 (1980)
- 8) Komaru A and Wada KT: Karyotypes of for species in the Pectinidae (Bivalvia: Pteriomorpha). *Bull Jap Jour Malac*, **44**, 249-259 (1985)
- 9) Komaru A and Wada KT: Electrophoretic survey of genetic variation in the scallop *Chlamys nobilis*. *Bull Nat Res Inst Aquaculture*, **9**, 7-13 (1986)
- 10) Komaru A and Wada KT: Gametogenesis and growth of induced triploid scallops *Chlamys nobilis*. *Nippon Suisan Gakkaishi*, **55**, 447-452 (1989)
- 11) Komaru A and Wada KT: Different processes of pronuclear events in pressure-treated and CB-treated zygotes at the second meiosis in scallop. *Nippon Suisan Gakkaishi*, **57**, 1219-1223 (1991)
- 12) Yamamoto K, Adachi S, Tamura I, Aramizu T and Koube H: Effects of hypoxia and water temperature on ciliary movement of gills 5 bivalvia, *Mytilus edulis*, *Atrina pectinate*, *Pinctada fucata martensii*, *Chlamys nobilis* and *Crassostrea gigas*. *J Nat Fish Univ*, **44**, 137-142 (1996)
- 13) Yamamoto K, Handa T, Araki A: The structure of the ctenidium in the noble scallop *Mimachlamys nobilis*. *J Nat Fish Univ*, **64**, 120-142 (2016)
- 14) Booth CE, McDonald DG, Walsh PJ: Acid-base balance in the sea mussel, *Mytilus edulis*. I. Effects of hypoxia and air-exposure on hemolymph acid-base status. *Mar Bio Lett*, **5**, 347-358 (1984)
- 15) Byrne RA, Shipman BN, Smatresk NJ, Dietz TH, McMahon RF : Acid-base balance during emergence in the freshwater bivalve *Corbicula fluminea*. *Physiol Zool*, **64**, 748-766 (1991)
- 16) Handa T, Yamamoto K: The blood acid-base balance in the pearl oyster, *Pinctada fucata martensii*, after the surgery. *J Nat Fish Univ*, **60**, 57-61 (2011)
- 17) Handa T, Yamamoto K: The acid-base balance of the hemolymph in the pearl oyster *Pinctada fucata martensii* under normoxic conditions. *Aquaculture Sci*, **60**, 113-117 (2012)
- 18) Handa T and Yamamoto K: Estimation of CO<sub>2</sub> partial pressure and bicarbonate concentration in the hemolymph of the black-lip pearl oyster *Pinctada margaritifera*. *J Nat Fish Univ*, **63**, 181-188 (2015)
- 19) Boutilier RG, Iwama GK, Heming TA, Randall DJ: The apparent pK of carbonic acid in rainbow trout blood plasma between 5 and 15 °C. *Resp. Physiol*, **61**, 237-254 (1985)
- 20) Yamamoto K, Handa T: Effect of hypoxia on oxygen uptake in the Pacific oyster *Crassostrea gigas*. *Aquaculture Sci*, **59**, 199-202, (2011)
- 21) Yamamoto K, Handa T, Nakamura M, Kitukawa K, Kita Y, Takimoto S, Nishikawa S: Effects of ozone-produced oxidants on respiration of the pearl oyster, *Pinctada fucata martensii*. *Aquaculture Sci*, **47**, 241-248 (1999)
- 22) Davenport HW: Fundamental equation. In: The ABC of acid-base chemistry 6th edition. University of Chicago Press, Chicago, 39-41 (1974)
- 23) Byrne AR, McMahon RB: Acid-base and ion regulation, during and following emersion, in the freshwater bivalve *Anodonta grandis simpsoniana*. *Biol Bull*, **181**, 289-297 (1991)
- 24) Michaelidis B, Haas D, Grieshaber MK: Extracellular and intracellular acid-base status with regard to the energy metabolism in the oyster *Crassostrea gigas* during exposure to air. *Physiol Biochem Zool*, **78**, 373-383 (2005a)
- 25) Michaelidis B, Ouzounis C, Paleras A, Portner HO: Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Mar Ecol Prog Ser*, **293**, 109-118 (2005b)
- 26) Cameron JN: The solubility of carbon dioxide as a function of temperature and salinity (Appendix table): In: Cameron JN (ed) Principles of physiological measurement. Academic Press, United Kingdom, 258-259 (1986)
- 27) Lindinger MI, Lauren DJ, McDonald DG: Acid-base balance in the sea mussel, *Mytilus edulis*. III. Effects of environmental hypercapnia on intra- and extracellular acid-base balance. *Mar Bio Lett*, **5**, 371-381 (1984)
- 28) Thomas RC: Intracellular pH. In: Hainsworth R (ed) Acid-base balance. Manchester University Press, United Kingdom, 50-74 (1986)
- 29) Truchot JP: Acid-base changes on transfer between seawater and freshwater in the Chinese crab *Eriocheir sinensis*. *Resp Physiol*, **87**, 419-427 (1992)
- 30) Whiteley MN, Scott, LJ, Breeze JS, Mccann L: Effects of water salinity acid-base balance in decapod crustaceans. *J Exp Biol*, **204**, 1003-1011 (2001)
- 31) Taylor WE, Whiteley MN: Oxygen transport and acid-base balance in the haemolymph of the lobster *Homarus gammarus* during aerial exposure and resubmersion. *J exp*

*Biol*, **144**, 417-436 (1989)

- 32) Rose AR, Wilkens LJ, Walker LR: The effects of walking on the heart rate, ventilation rate and acid-base status in the lobster *Homarus americanus*. *J Exp Biol*, **201**, 2601-2608 (1998)
- 33) Azuma T, Itazawa Y: Respiration of a marine teleost, the porgy, under normoxic, resting condition. *Nippon Suisan Gakkaishi*, **59**, 621-626 (1993)
- 34) Wood MC, McDonald GD, McMahon RB: The influence of experimental anaemia on blood acid-base regulation in vivo and in vitro in the starry flounder *Platichthys stellatus* and the rainbow trout *Salmo gairdneri*. *J exp Biol*, **96**, 221-237 (1982)
- 35) Nelson AJ, Tang Y, Boutilier GR; The effects of salinity change on the exercise performance of two Atlantic cod (*Gadus morhua*) populations inhabiting different environments. *J Exp Biol*, **199**, 1295-1309 (1996)
- 36) Itazawa Y, Takeda T: Gas exchange in the carp gills in normoxic and hypoxic conditions. *Resp Physiol*, **35**, 263-269 (1978)
- 37) Handa T, Yamamoto K: Blood and cardiovascular-respiratory properties of the common carp *Cyprinus carpio* under resting conditions. *J Nat Fish Univ*, **62**, 19-29 (2013)