

2014年山口県阿武町奈古定置網の伝票をもとにした 水揚げ個体数の推定に関する予備的調査

メタデータ	言語: English 出版者: 水産大学校 公開日: 2024-10-11 キーワード (Ja): キーワード (En): Catch slips; Fish catches; Fisheries; Set net 作成者: 田上, 英明, 井町, 博明, 水津, 辰則, 横田, 峻, 永井, 節子, 福村, 直人, 梶川, 和武, 毛利, 雅彦 メールアドレス: 所属: 水産研究・教育機構, 水産研究・教育機構, 水産研究・教育機構
URL	https://fra.repo.nii.ac.jp/records/2012132

This work is licensed under a Creative Commons
Attribution 4.0 International License.



Preliminary Study on Estimation of Abundance based on the Slips regarding the Catches Landed by Set Net off Nago, Abu Town, Yamaguchi Prefecture, Japan in 2014

Hideaki Tanoue^{1†}, Hiroaki Imachi², Takanori Suizu³, Shun Yokota²,
Setsuko Nagai³, Naoto Fukumura³, Yoritake Kajikawa¹,
and Masahiko Mohri⁴

Abstract : Most of the slips for catches landed by set net fishery describe the number of boxes every catch shipped in Japan. The number of the catches contained in each box (“*irisu*”) depends on the species. Furthermore, even among the same species, the “*irisu*” may change depending on the season. In this study, under the cooperation of fishermen, with reference to their catch slips, the abundance by species of the catches landed was estimated by researching the contents of the boxes in Nago, Abu Town, Yamaguchi Prefecture, Japan, 2014. The number of operation days was 206, which amounted to the monthly average number of operation days of 17.2 ± 3.4 (average $\pm$ SD). The number of species shipped in the same year amounted to 61 with the monthly average of 22.8 ± 5.8 . We interviewed fishermen to get the relationship between the species or fishes whose brand names may change depending on the season and the respective “*irisu*” . Based on the results of the interview, the total abundance of catches landed (exclude “*iromono*” boxes to be shipped with some different fish species) was estimated to 218,543 and the average abundance shipped per operation day amounted to 943.3 ± 1297.9 .

Key words : Catch slips, Fish catches, Fisheries, Set net

Introduction

Catch diversity of set net is one of the important items for understanding the characteristics of fishing conditions in coastal waters. There is one of the usual approaches of research the abundance data after purchasing all the catches obtained by set nets ¹⁾. On the other hand, Tanoue et al.²⁾ indicated the catches in set nets using diversity indexes in the previous paper, where there remained a subject of how to convert the number of boxes into the abundance and also they used an index which described the only species instead of the abundance data ³⁾.

Most of the slips for catches landed by set net fishery describe the number of boxes every catch shipped in

Japan. The number of the catches contained in each box (hereinafter referred to as “*irisu*”) depends on the species. Furthermore, even among the same species, the “*irisu*” may change depending on the season.

In this study, under the cooperation of fishermen involved in set net fishery, with reference to their catch slips, the abundance by species of the catches landed was estimated by researching the contents of the boxes to be offered on the market.

Materials and methods

This study uses the catch slip data from fishermen in Nago, Abu Town, Yamaguchi Prefecture. This area is located in the northern part of this prefecture along the

1. Department of Fisheries Science and Technology, National Fisheries University

2. Graduate student, Graduate School of Fisheries Science, National Fisheries University

3. Under graduate student, Department of Fisheries Science and Technology, National Fisheries University

4. Graduate School of Fisheries Science, National Fisheries University

† Hideaki Tanoue : h-tanoue@fish-u.ac.jp

Sea of Japan (34°30' 11.9" N, 131°28' 17" E). The catch data are based on the species and the number of boxes filled in the slip for 2014. Usually, the figures indicated on the slips stand for the number of boxes. In this study, we interviewed fishermen to get the relationship between the species or fishes whose brand names may change depending on the season and the respective "irisū". Based on the results of the interview, the abundance was estimated. Scientific names and identification of species follow Nakabo ⁴⁾ and Okutani ⁵⁾.

Results and Discussion

The number of operation days of set net fishery in the Nago area of Abu Town in 2014 was 206 (Table 1), which amounted to the monthly average number of operation days of 17.2 ± 3.4 (average $\pm$ SD) (Max: 23 days in February, Min: 12 days in September). The number of species shipped in the same year amounted to 61 with the monthly average of 22.8 ± 5.8 .

The results of the interview showed as follows: For fish species, the figure filled in the "brand names" column corresponded to the "irisū". However, for squid species, the figure filled in the same column did not correspond to the "irisū" (Table 2), which means that different methods of calculating the number of individuals must be applied to fishes and squids. Furthermore, spear squid (*Heterololigo bleekeri*) and swordtip squid (*Uroteuthis edulis*) were caught in nets, some transactions ("Katsugyo": keep their alive in tank) were made on a weight basis. In such a case, the transaction was made in units of one kg and it was found that approximately three squids weighed one kg. Therefore, the individual data based on the weight should be calculated by multiplying the total weight (kg)/day by 3. Especially, extremely small size of squids and fishes (*Trachurus japonicus*) were shipped approximately 5 kg/box, which is called "baraika" and "zengoaji" in the Nago area. We calculated that the number of the catches contained in the box by dividing 5 kg by average of individual weight according to random sampling.

On the other hand, there is a box to be shipped with

some different fish species because the catch does not amount to the "irisū" of a certain brand, which is called "iromono". Such a box was identified for 133 (64.4%) out of 206 operation days, which accounted for 3.8 ± 6.7 % of the total amount of money of the catches landed per day. In this study, the abundance of such a box is not considered.

Based on the results of the interview, the calculation was completed by adding the individual data estimated by multiplying the number of fish boxes by the "irisū" to that estimated from the weights of the squids and fishes. The calculation results showed that the total abundance of catches landed was estimated to 218,543 and the average abundance shipped per operation day amounted to 943.3 ± 1297.9 (Table 3).

Acknowledgement

We thank the Mr.H.Suizu of the Nago Branch of the Yamaguchi Prefectural Fisheries Cooperative Association. This study was financially supported by Hagi city and JSPS KAKENHI Grant Number 15K18739.

References

- 1) Michine A, Miura T, Sasaki T: Characteristics in occurrence of fish and crustacean collected with set net in Lake Nakaumi, estuarine inland-sea, western Japan, *Rep Shimane Pref Fish Technol Cent*, **5**, 53-65 (2013) (in Japanese)
- 2) Tanoue H, Nozoe A, Fujiwara K, Mohri M: Changes in Fishes Caught by Set Net Fishery Observed by Taxonomic Distinctness: Preliminary Study Using the Set Net Data on Futaoi Island, Simonoseki, Japan, *Math Phys Fish Sci*, **13**, 1-9 (2016)
- 3) Tanoue H, Imachi H, Nozoe A, Mohri M: Taxonomic Diversity ($\Delta +$) of Fishes Caught by Set Net Fishery on Futaoi-jima Island, Shimonoseki, Japan, *J Nat Fish Univ*, **65**, 261-266 (2017) (in Japanese)
- 4) Nakabo T (ed). Fishes of Japan with pictorial keys to the species, third edition. Tokai University Press, Hadano (2013) (in Japanese)

- 5) Okutani T: Cuttlefishes and squids of the world [New Edition], Tokai University Press, Hadano (2015) (in Japanese)

Table 1. Summary of operation-days and landing by set net off Abu town in 2014

Operation days		No. of species		Landed quantity (abundance)	
Total	days / month ($\pm$ SD)	Total	species / month ($\pm$ SD)	Total	individuals / day ($\pm$ SD)
206	17.2 $\pm$ 3.4	61	22.8 $\pm$ 5.8	218,543	943.3 $\pm$ 1297.9

Table 2. Relationship between catch slips of brand names (code number) and the number of squids contained (squid's "irisu") in box

Species	Brand names (Code number)											
	Uncoded	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0
Spear squid (<i>Heterololigo bleekeri</i>)	14	-	-	-	17	27(24)	-	-	-	-	-	-
Swordtip squid (<i>Uroteuthis edulis</i>)	14	-	-	-	17	24	28	36	40	50	55	66
Bigfin reef squid (<i>Sepioteuthis lessoniana</i>)	2	-	3	4	6(3)	12	16(14)	20(16)	22	25	-	-
Golden cuttlefish (<i>Sepia esculenta</i>)	8	-	-	-	-	-	-	-	-	-	-	-
Japanese flying squid (<i>Todarodes pacificus</i>)	14	-	-	-	-	-	-	-	-	-	-	-
Rhomboid squid (<i>Thysanoteuthis rhombus</i>)	1	-	-	-	-	-	-	-	-	-	-	-

Number shown in parentheses is "irisu", especially from December to January in winter.

Table 3. Results of landed species and quantity by set net off Abu town in 2014

		Species	Landed quantity (n)
Cephalopoda	Sepiida	Sepiidae	<i>Sepia esculenta</i> 152
	Myopsida	Loliginidae	<i>Uroteuthis edulis</i> 143,454
			<i>Heterololigo bleekeri</i> 37,220
			<i>Sepioteuthis lessoniana</i> 2,978
	Teuthida	Ommastrephidae	<i>Todarodes pacificus</i> 2,282
		Thysanoteuthidae	<i>Thysanoteuthis rhombus</i> 9
	Octopoda		
		Octopodidae	<i>Octopus vulgaris</i> 19
Chondrichthyes	Carcharhiniformes		
		Triakidae	<i>Triakis scyllium</i> 5
	Squatiniformes		
		Squatinidae	<i>Squatina japonica</i> 9
	Rhinobatiformes		
		Rhinobatidae	<i>Rhinobatos schlegelii</i> 12
	Myliobatiformes	Dasyatidae	<i>Dasyatis akajei</i> 6
Osteichthyes	Anguilliformes		
		Congridae	<i>Conger myriaster</i> 2
	Salmoniformes		
		Slamonidae	<i>Oncorhynchus keta</i> 17
			<i>Oncorhynchus masou masou</i> 2
	Clupeiformes		
		Clupeidae	<i>Konosirus punctatus</i> 2
	Gadiformes		
		Gadidae	<i>Gadus macrocephalus</i> 5
	Zeiformes		
		Zeidae	<i>Zeus faber</i> 77
	Gasterosteiformes		
		Fistulariidae	<i>Fistularia petimba</i> 6
	Perciformes		
		Sebastidae	<i>Sebastes spp.</i> 4
			<i>Sebastes marmoratus</i> 9
		Synanceiida	<i>Inimicus japonicus</i> 15
		Triglidae	<i>Chelidonichthys spinosus</i> 33
		Lateolabracidae	<i>Lateolabrax japonicus</i> 1,072

	<i>Lateolabrax latius</i>	45
Serranidae	<i>Epinephelus bruneus</i>	4
	<i>Epinephelus akaara</i>	16
Coryphaenidae	<i>Coryphaena hippurus</i>	18
Carangidae	<i>Seriola quinqueradiata</i>	8,824
	<i>Seriola aureovittata</i>	1,570
	<i>Seriola dumerili</i>	107
	<i>Trachurus japonicus</i>	6,762
Bramidae	<i>Brama japonica</i>	1
Haemulidae	<i>Parapristipoma trilineatum</i>	31
	<i>Diagramma picta</i>	11
Sparidae	<i>Acanthopagrus schlegelii</i>	16
	<i>Pagrus major</i>	105
	<i>Rhabdosargus sarba</i>	1
Girellidae	<i>Girella punctata</i>	1,672
Kyphosidae	<i>Kyphosus vaigiensis</i>	6
Oplegnathidae	<i>Oplegnathus punctatus</i>	56
	<i>Oplegnathus fasciatus</i>	32
Cheilodactylidae	<i>Goniistius zonatus</i>	2
Siganidae	<i>Siganus fuscescens</i>	146
Acanthuridae	<i>Prionurus scalprum</i>	7
Trichiuridae	<i>Trichiurus japonicus</i>	1
Scombridae	<i>Scomber japonicus</i>	2,030
	<i>Acanthocybium solandri</i>	1
	<i>Scomberomorus niphonius</i>	2,479
	<i>Thunnus orientalis</i>	25
	<i>Thunnus tonggol</i>	15
Istiophoridae	<i>Istiophorus platypterus</i>	1
Pleuronectiformes		
Paralichthyidae	<i>Paralichthys olivaceus</i>	206
Pleuronectidae	<i>Pleuronectes herzensteini</i>	18
Tetraodontiformes		
Monacanthidae	<i>Stephanolepis cirrhifer</i>	26
	<i>Aluterus Monoceros</i>	92
Tetraodontidae	<i>Takifugu pordalis</i>	2,900
	<i>Takifugu xanthopterus</i>	3
	<i>Lagocephalus spadiceus</i>	45
	<i>Takifugu rubripes</i>	29
	<i>Takifugu flavipterus</i>	3,850

2014年山口県阿武町奈古定置網の伝票をもとにした 水揚げ個体数の推定に関する予備的調査

田上英明^{1†}・井町博明²・水津辰則³・横田峻²・永井節子³・
福村直人³・梶川和武¹・毛利雅彦⁴

要 旨

定置網の水揚げ伝票には、出荷された漁獲物が箱数で記載されている場合が多い。箱に入っている漁獲物の個体数（以下、入数）は種によって異なり、同じ種であったとしても季節によって入数に変化する場合がある。本研究では、山口県阿武町奈古の定置網で出荷される漁獲物の個体数を推定するために、漁業者に協力いただき、2014年の種また季節で変化する可能性がある入数を聞き取り調査によって調べることを検討した。2014年の阿武町奈古の定置網の操

業日数は206日、月当たりの平均操業日数は 17.2 ± 3.4 （平均 $\pm$ SD）日であった。出荷種数は61種、月平均は 22.8 ± 5.8 種であった。聞き取り調査の結果をもとに市場に揚がる箱の内容物を確認し、水揚げされた種毎の個体数を推定した。その結果、入数に数が達せず、複数の魚種を混ぜ出荷された箱（イロもの、1日辺り水揚げ金額の $3.8 \pm 6.7\%$ ）を除いた2014年の水揚げ総個体数は、218,543個体と推定され、操業日当たりの推定平均出荷個体数は 943.3 ± 1297.9 個体であった。

1. 水産大学校海洋生産管理学科 (Department of Fisheries Science and Technology, National Fisheries University)
2. 水産大学校水産学研究科学生 (Graduate student, Graduate School of Fisheries Science, National Fisheries University)
3. 水産大学校海洋生産管理学科学生 (Under graduate student, Department of Fisheries Science and Technology, National Fisheries University)
4. 水産学研究科 (Graduate School of Fisheries Science, National Fisheries University)