

## Effect of Air Exposure on the Acid-Base Balance of Hemolymph in Black-lip Pearl Oyster *Pinctada margaritifera*

メタデータ	言語: English 出版者: 水産大学校 公開日: 2024-10-11 キーワード (Ja): キーワード (En): <i>Pinctada margaritifera</i> ; black-lip pearl oyster; hemolymph; acid?base balance; air exposure; respiratory physiology 作成者: 半田, 岳志, 荒木, 晶, 山元, 憲一 メールアドレス: 所属: 水産研究・教育機構
URL	<a href="https://fra.repo.nii.ac.jp/records/2012166">https://fra.repo.nii.ac.jp/records/2012166</a>

This work is licensed under a Creative Commons Attribution 4.0 International License.



# Effect of Air Exposure on the Acid-Base Balance of Hemolymph in Black-lip Pearl Oyster *Pinctada margaritifera*

HANDA Takeshi,<sup>1†</sup> ARAKI Akira<sup>1</sup> and YAMAMOTO Ken-ichi<sup>1</sup>

**Abstract** : We investigated the hemolymph acid-base status of the black-lip pearl oyster *Pinctada margaritifera* exposed to air at 23°C. Air-exposed *P. margaritifera* showed a decrease in hemolymph pH from 7.563 to 6.965 after 36 h. The hemolymph total CO<sub>2</sub> concentration (Tco<sub>2</sub>) increased from 2.01 mM/L to 5.63 mM/L during 36 h of air exposure. The hemolymph CO<sub>2</sub> partial pressure (Pco<sub>2</sub>) and bicarbonate concentration ([HCO<sub>3</sub><sup>-</sup>]) were calculated, by rearranging the Henderson-Hasselbalch equation with substitution of CO<sub>2</sub> solubility coefficient ( $\alpha_{\text{CO}_2}$ , 40.49 mM/L/torr) and apparent dissociation constant of carbonic acid (pKapp, 5.98351). The hemolymph Pco<sub>2</sub> increased from 1.28 torr to 13.68 torr, and [HCO<sub>3</sub><sup>-</sup>] increased from 1.96 mM/L to 5.08 mM/L during 36 h of air exposure. The hemolymph calcium concentration ([Ca<sup>2+</sup>]) increased from 7.96 mM/L to 11.2 mM/L. These results indicated that during prolonged air exposure *P. margaritifera* showed hemolymph acidosis with partial metabolic compensation by the mobilized bicarbonate from the shell valve. When the experimental animals were immersed in seawater after air exposure, *P. margaritifera* needed 24 hours at least for recovery of the hemolymph acid-base status to the initial level.

**Key words** : *Pinctada margaritifera*, black-lip pearl oyster, hemolymph, acid-base balance, air exposure, respiratory physiology

## Introduction

The black-lip pearl oyster *Pinctada margaritifera* is a filibranchial bivalve classified in the Pteriidae<sup>1)</sup> and is distributed in the Indo-West Pacific region near the equator, including in the Red Sea, Arabian Sea, Persian Gulf, and around India, Sri Lanka, New Guinea, Hawaii and Madagascar.<sup>2)</sup> In Japan, *P. margaritifera* is present southwards from the Kii Peninsula.<sup>1)</sup> The black-lip pearl oyster has nacreous aragonite in the inner layer of its shell valves, and it is used for black pearl production. The process of pearl production is similar to the growth of the shell and is related directly to metabolism. The metabolism of the black-lip pearl oyster has been studied in terms of regulation of gill ventilation volume and oxygen uptake in normoxic, hypoxic, and anathermal conditions.<sup>3-6)</sup> The anatomical structures of the ctenidium, circulatory system, digestive diverticula, and labial palp have been studied.<sup>7,8)</sup> Handa et al. (2015) described the estimation of hemolymph CO<sub>2</sub> partial pressure (Pco<sub>2</sub>) of

black-lip pearl oyster in normoxic conditions.<sup>9)</sup> However, there are few reports of the effect of air exposure on the respiratory physiology from the viewpoint of CO<sub>2</sub> dynamics phase and acid-base balance in black-lip pearl oyster. In the pearl production, pearl oysters are often exposed to the air for the surgical implantation of a nucleus. The host pearl oysters are removed from the seawater, and a wedge is placed between the shell valves to keep the pearl oyster open.<sup>10)</sup> The oyster is clamped open, placed in the shell stand, and undergoes a surgical operation.<sup>10)</sup> Therefore, research into the effect of air exposure on the acid-base balance may contribute to efficient CO<sub>2</sub> utilization and growth, which are related to pearl formation and the handling of black-lip pearl oysters. We examined the hemolymph acid-base status of the black-lip pearl oyster *P. margaritifera* pre- and post-air exposure, and evaluated the acid-base balance and CO<sub>2</sub> dynamics *in vivo*. The hemolymph Pco<sub>2</sub> values of marine bivalves, including *P. margaritifera*, are very low

Affiliation:<sup>1</sup> Department of Applied Aquabiology, National Fisheries University, Nagata-honmachi, Shimonoseki, Yamaguchi, JAPAN  
† Corresponding author: handat@fish-u.ac.jp (T. HANDA)

(0.57–2.3 torr) in normoxic and normocapnic conditions.<sup>9,11-19</sup> The estimation of  $P_{\text{CO}_2}$  by application of the Henderson–Hasselbalch equation is practiced in studies of the acid–base balance owing to its relative ease and accuracy.<sup>20</sup> In the equation, the  $\text{CO}_2$  solubility coefficient ( $\alpha_{\text{CO}_2}$ ) and apparent dissociation constant (pKapp) of carbonic acid in the hemolymph are required for the experimental animal. Thus, we determined the coefficient and constant of the hemolymph *in vitro*, and calculated the hemolymph  $P_{\text{CO}_2}$  of *P. margaritifera*, and available for the evaluation of the hemolymph acid–base balance.

## Materials and Methods

### Experimental animals and conditions

The experiments used 29 black-lip pearl oysters (mean total wet weight: 188 g). After cleaning the shell valves, they were reared for one month at 23°C in aerated seawater with added cultivated phytoplankton.<sup>21-23</sup> Twenty-four hours before collecting hemolymph, the black-lip pearl oysters were transferred to a respiratory chamber with a flow of particle-free (>0.45  $\mu\text{m}$ ) seawater. All experiments were conducted in seawater with a salinity of 33 psu, water temperature 23°C,  $\text{O}_2$  saturation 98%, pH 8.15, and total  $\text{CO}_2$  content 1.5 mM/L.

### Experimental procedure

The effect of air exposure on hemolymph acid–base status was investigated *in vivo*, and hemolymph  $P_{\text{CO}_2}$  and bicarbonate concentration ( $[\text{HCO}_3^-]$ ) were calculated using the results of *in vitro* experiments in this study.

#### *In vivo* experiments

Experimental animals in the respiratory chamber were exposed to air for 36 h by stopping the flow into the chamber and siphoning out the seawater. The temperature and humidity of the air were maintained by passing air through the experimental seawater, and adjusted air flowed into the respiratory chamber. Hemolymph was collected from the adductor muscle of the air-exposed animals at 36 h (AE36h). The inflow of

experimental seawater was resumed into the respiratory chamber after exposure to air for 36 h, and the animals were immersed in seawater. The hemolymph of immersed animals was collected at 6 h (Im6h) and 24 h (Im24h). As a control for the experimental animals, hemolymph was collected before air exposure (AE0h). Six different individual animals were used for each analysis (n=24).

#### *In vitro* experiments

The hemolymph  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  were calculated by rearranging the Henderson–Hasselbalch equation.<sup>20,24</sup> In the equation, the  $\alpha_{\text{CO}_2}$  and pKapp were required in the black-lip pearl oyster. *In vitro* determination of  $\alpha_{\text{CO}_2}$  and pKapp were performed on hemolymph drawn from the adductor muscle of the experimental animals before exposure to air (n=5).  $\alpha_{\text{CO}_2}$  was determined using hemolymph, which was adjusted to pH 2.5 by the addition of lactic acid (Wako Pure Chemical Co., JP). The acidified sample was transferred to a tonometer flask and equilibrated with humidified standard  $\text{CO}_2$  gas ( $\text{CO}_2$ , 15.0%;  $\text{O}_2$ , 20.9%;  $\text{N}_2$  Balance) using an equilibrator (DEQ-1, Cameron Instruments) at 23°C, and subsequently the total  $\text{CO}_2$  concentration ( $T_{\text{CO}_2}$ , mM/L) of each equilibrated sample was measured. The  $P_{\text{CO}_2}$  (torr) of the equilibrated sample was calculated from known  $\text{CO}_2$  concentration standard gas (15.0%), prevailing barometric pressure and water vapor pressure at 23°C. For the determination of pKapp, the hemolymph sample was transferred to a tonometer flask and equilibrated with humidified standard  $\text{CO}_2$  gases ( $\text{CO}_2$ , 0.2, 0.5, 1.0, and 2.0%;  $\text{O}_2$ , 20.9%;  $\text{N}_2$  Balance) using an equilibrator at 23°C. After equilibration, the pH and  $T_{\text{CO}_2}$  of the sample were measured. Using the sample pH,  $T_{\text{CO}_2}$ , and  $\alpha_{\text{CO}_2}$ , the pKapp was determined by rearrangement of the Henderson–Hasselbalch equation.

#### Hemolymph collection and analysis

Hemolymph was collected once from each individual from the adductor muscle by direct puncture with a gas-tight microsyringe (Model 1750LTN, Hamilton Co., US), and the volume of each hemolymph sample was 0.4–0.5

mL. Hemolymph pH and  $Tco_2$  were measured immediately after collection. The pH was measured using a blood gas meter (BGM200; Cameron Instruments Co., US) with glass and reference electrodes (E301, E351; Cameron Instruments Co., US) at 23°C.  $Tco_2$  was measured using a total  $CO_2$  analyzer (Capnicon 5; Cameron Instruments Co., US). Hemolymph calcium concentrations ( $[Ca^{2+}]$ , mM/L) were determined with a test kit (Calcium E-test, Wako Pure Chemical Co., JP) and a spectrophotometer (Spectronic 20A, Shimadzu Co., JP).

### Calculation

$aco_2$  of the experimental animals was calculated using the equation:

$$aco_2 = Tco_2 \cdot Pco_2^{-1}$$

where the units of the parameters are mM/L/torr in  $aco_2$ , mM/L in  $Tco_2$  and torr in  $Pco_2$ .

Using the sample pH,  $Tco_2$ , and  $aco_2$  calculated using the above equation,  $pK_{app}$  was determined by rearrangement of Henderson–Hasselbalch equation<sup>20,24</sup> as follows:

$$pK_{app} = pH - \log [(Tco_2 - aco_2 \cdot Pco_2) \cdot (aco_2 \cdot Pco_2)^{-1}]$$

where  $Pco_2$  is calculated from known  $CO_2$  concentration standard gases.

The hemolymph  $Pco_2$  and  $[HCO_3^-]$  (mM/L) were calculated using the equations:

$$Pco_2 = Tco_2 \cdot [aco_2 \cdot (1 + 10^{(pH - pK_{app})})]^{-1}$$

$$[HCO_3^-] = Tco_2 - aco_2 \cdot Pco_2$$

where  $Tco_2$  and pH were measured values, and  $aco_2$  and  $pK_{app}$  were obtained in *in vitro* experiments.

The non-bicarbonate buffer value ( $\beta_{NB}$ , slykes) and the relational expression of the hemolymph non-bicarbonate buffer were calculated from pH and  $[HCO_3^-]$  of the *in vitro* experiment. Apparent buffer value of the *in vivo* experiment ( $\beta_{vivo}$ , slykes) was calculated from the hemolymph pH and  $[HCO_3^-]$  of the animals exposed to air


at AE0h and AE36h.

### Statistical analysis

Data are expressed as means  $\pm$  standard error of the means. Kruskal–Wallis test was performed for changes in hemolymph properties over the experimental time course. The multiple comparison of all pairs used the Steel–Dwass test. Statistically significant differences were set at  $P < 0.05$  (KyPlot 5.0, KyensLab Inc., JP).

## Results

The mean values of hemolymph pH were statistically significantly decreased from 7.563 to 6.965 during air exposure for 36 h ( $P < 0.05$ , Fig. 1). The hemolymph  $Tco_2$  increased from 2.01 mM/L to 5.63 mM/L during air exposure for 36 h ( $P < 0.05$ , Fig. 2). When the experimental animals were immersed in the seawater after air exposure, the hemolymph pH increased and  $Tco_2$  decreased at Im6h and Im24h ( $P < 0.05$ ). There were no significant differences in hemolymph pH and  $Tco_2$  between control and immersed animals (Figs. 1-2). In *in vitro* experiments, hemolymph  $aco_2$  was 40.49  $\mu$ M/L/torr. The hemolymph  $pK_{app}$  at known  $Pco_2$  (standard gases)




**Fig. 1** Effect of air exposure on hemolymph pH in black-lip pearl oyster *Pinctada margaritifera*. AE0h: control; AE36h: air exposure for 36 h; Im6h: immersion for 6 h after air exposure; Im24h: immersion for 24 h after air exposure. Hemolymph was collected from each experimental animal (n=6 in each column). Values are means  $\pm$  SEM. Different lowercase letters indicate statistically significant differences ( $P < 0.05$ , Steel–Dwass multiple comparison test).

and the corresponding measured pH and  $T_{\text{CO}_2}$  are shown in Table 1. The calculated  $pK_{\text{app}}$  from all hemolymph samples was  $5.98351 \pm 0.014536$ .  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  were calculated by substitution of the values of  $a_{\text{CO}_2}$  and  $pK_{\text{app}}$  in the rearranged Henderson–Hasselbalch equation as follows:

$$P_{\text{CO}_2} = T_{\text{CO}_2} \cdot [0.04049 \cdot (1 + 10^{(\text{pH} - 5.98351)})]^{-1}$$


$$[\text{HCO}_3^-] = T_{\text{CO}_2} - 0.04049 \cdot P_{\text{CO}_2}$$

where the units of the parameters are torr for  $P_{\text{CO}_2}$  and mM/L for  $T_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$ .



**Fig. 2** Effect of air exposure on hemolymph total  $\text{CO}_2$  concentration ( $T_{\text{CO}_2}$ ) in black-lip pearl oyster *Pinctada margaritifera*. AE0h: control; AE36h: air exposure for 36 h; Im6h: immersion for 6 h after air exposure; Im24h: immersion for 24 h after air exposure. Hemolymph was collected from each experimental animal ( $n=6$  in each column). Values are means  $\pm$  SEM. Different lowercase letters indicate statistically significant differences ( $P<0.05$ , Steel–Dwass multiple comparison test).

The mean values of the hemolymph  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  of the control values (AE0h) were calculated as 1.28 torr and 1.96 mM/L, respectively (Figs. 3, 4). The hemolymph  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  increased significantly during air exposure ( $P<0.05$ ), reaching 13.6 torr and 5.08 mM/L at AE36h. When air-exposed animals were immersed in seawater, hemolymph  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  decreased at Im6h and Im24h. There were no significant differences in hemolymph  $P_{\text{CO}_2}$  and  $[\text{HCO}_3^-]$  between control and immersed animals. Hemolymph  $[\text{Ca}^{2+}]$  increased statistically significantly from 7.96 mM/L to 11.2 mM/L during air exposure for 36 h ( $P<0.05$ , Fig. 5). In immersed




**Fig. 3** Effect of air exposure on hemolymph  $\text{CO}_2$  partial pressure ( $P_{\text{CO}_2}$ ) in black-lip pearl oyster *Pinctada margaritifera*. AE0h: control; AE36h: air exposure for 36 h; Im6h: immersion for 6 h after air exposure; Im24h: immersion for 24 h after air exposure. Hemolymph was collected from each experimental animal ( $n=6$  in each column). Values are means  $\pm$  SEM. Different lowercase letters indicate statistically significant differences ( $P<0.05$ , Steel–Dwass multiple comparison test).

**Table 1.** Mean values of hemolymph pH, bicarbonate ion concentration ( $[\text{HCO}_3^-]$ ) and apparent dissociation constant of carbonic acid ( $pK_{\text{app}}$ ) of the black-lip pearl oyster (*Pinctada margaritifera*) with known  $\text{CO}_2$  standard gases at 23°C


Standard gas		Hemolymph			
$\text{CO}_2$ (%)	$P_{\text{CO}_2}$ (torr)	pH	$[\text{HCO}_3^-]$ (mM/L)	$pK_{\text{app}}$	N
0.203	1.50	7.360	1.86	5.87958	5
0.515	3.81	7.119	1.94	6.02382	5
1.01	7.46	6.839	1.97	6.02678	5
2.00	14.8	6.589	2.32	6.00384	5

Barometric pressure, 767 $\pm$ 4.1 torr;  $\text{CO}_2$  solubility coefficient ( $a_{\text{CO}_2}$ ), 40.49  $\mu\text{M/L/torr}$

animals, hemolymph  $[Ca^{2+}]$  at Im6h (10.4 mM/L) was statistically significantly higher than the control value (AE0h, 7.96 mM/L), although hemolymph  $[Ca^{2+}]$  at Im24h (8.94 mM/L) was approximately the same as the control value. The progress of change in acid-base balance in experimental animals is summarized in a pH- $[HCO_3^-]$



**Fig. 4** Effect of air exposure on hemolymph bicarbonate concentration ( $[HCO_3^-]$ ) in black-lip pearl oyster *Pinctada margaritifera*. AE0h: control; AE36h: air exposure for 36 h; Im6h: immersion for 6 h after air exposure; Im24h: immersion for 24 h after air exposure. Hemolymph was collected from each experimental animal ( $n=6$  in each column). Values are means  $\pm$  SEM. Different lowercase letters indicate statistically significant differences ( $P<0.05$ , Steel-Dwass multiple comparison test).




**Fig. 5** Effect of air exposure on hemolymph calcium ion concentration ( $[Ca^{2+}]$ ) in black-lip pearl oyster *Pinctada margaritifera*. AE0h: control; AE36h: air exposure for 36 h; Im6h: immersion for 6 h after air exposure; Im24h: immersion for 24 h after air exposure. Hemolymph was collected from each experimental animal ( $n=6$  in each column). Values are means  $\pm$  SEM. Different lowercase letters indicate statistically significant differences ( $P<0.05$ , Steel-Dwass multiple comparison test).

diagram (Fig. 6). The hemolymph  $[HCO_3^-]$  of air-exposed animals rose with decreasing pH, and the point at AE36h was located above the non-bicarbonate buffer line, for which the relationship with the expression is indicated in Fig. 6. The hemolymph pH and  $[HCO_3^-]$  of immersed animals were similar to the control values, with the points located near the non-bicarbonate buffer line. The  $\beta_{NB}$ , which is the slope of the non-bicarbonate buffer line, was 0.53 slykes. The  $\beta_{inv}$ , which was calculated from the hemolymph pH and  $[HCO_3^-]$  of air-exposed animals at AE0h and AE36h, was 5.22 slykes.

## Discussion

We examined the hemolymph acid-base status of the black-lip pearl oyster *P. margaritifera* to evaluate the effect of air exposure on acid-base balance and  $CO_2$  dynamics phase. *P. margaritifera* showed a reduction in hemolymph pH and increases in  $Tco_2$  and  $Pco_2$  during air exposure for 36 h. The air-exposed animals were unable to ventilate the gill, which inhibited the release of  $CO_2$ .  $CO_2$  gradually accumulated in the hemolymph, causing progressive acidosis. Therefore, *P. margaritifera* during



**Fig. 6** Hemolymph pH- $[HCO_3^-]$  diagram of air-exposed (closed circles), immersed (open squares), and control (open circles) black-lip pearl oysters *Pinctada margaritifera*. The  $Pco_2$  isopleths are derived from rearranging the Henderson-Hasselbalch equation,<sup>24)</sup> and  $a_{CO_2}$  and  $pK_{app}$  of the experimental animals were 40.49  $\mu M/L/torr$  and 5.98351, respectively. The dashed line is the non-bicarbonate buffer line:  $[HCO_3^-] = 5.7337 - 0.5319 \cdot pH$  ( $R^2=0.7858$ ). The line was obtained from the results of *in vitro* experiments in this study.

the air exposure should include respiratory acidosis due to the inhibition of CO<sub>2</sub> release. In some marine and freshwater bivalves, the hemolymph and pericardiac fluid showed a drop of oxygen partial pressure<sup>11,25,26)</sup> and acidosis during air exposure.<sup>11,25-29)</sup> Although we did not measure the anaerobic end-products, the results of biochemical studies on anaerobic metabolism<sup>30-34)</sup> suggested that air exposure in this study was sufficient to force anaerobic metabolism. *P. margaritifera* exposed to air for a long time might undergo metabolic acidosis due to anaerobic metabolism. *P. margaritifera* exposed to the air increased hemolymph [HCO<sub>3</sub><sup>-</sup>] and [Ca<sup>2+</sup>] at AE36h. The increased [HCO<sub>3</sub><sup>-</sup>] and [Ca<sup>2+</sup>] during air exposure seemed to be mobilized from CaCO<sub>3</sub> crystals in the shell valves in *P. margaritifera*. In marine and freshwater bivalves, acidosis during air exposure induces increases in [HCO<sub>3</sub><sup>-</sup>] and [Ca<sup>2+</sup>] in the hemolymph or pericardiac fluid.<sup>11,25-27)</sup> Research using radiolabeled markers indicated that the source of increased calcium was the shell valve.<sup>35)</sup> The increase in acidic end-products of anaerobic metabolism could dissolve the shell valve of *P. margaritifera*, and the bicarbonate and calcium ions were mobilized from the shell valves to the hemolymph during air exposure in this study. The mobilized bicarbonate seemed to be effective for buffering acidosis in *P. margaritifera* hemolymph.

When the experimental animals were immersed in seawater, *P. margaritifera* increased hemolymph pH at Im6h. The hemolymph Pco<sub>2</sub> was reduced with the increase in pH and the decrease in [HCO<sub>3</sub><sup>-</sup>]. The immersed animals should resume gill ventilation and rapidly release CO<sub>2</sub> from the gill into the seawater for 6 h. The immersed animals might release CO<sub>2</sub> by the diffusion from the surface of the soft body. Aerobic metabolism resumed in the immersed animals, and the production of anaerobic acidic end-products stopped. The increased [HCO<sub>3</sub><sup>-</sup>] during air exposure was consumed to compensate for the acidosis within 6 h in the immersed animals, and [HCO<sub>3</sub><sup>-</sup>] decreased to the control level. On the other hand, the increased [Ca<sup>2+</sup>] during air exposure did not return to the control level within 6 h in immersed animals, although a declining trend in hemolymph [Ca<sup>2+</sup>]

was shown. Silverman et al. (1983) reported that the freshwater mussel *Ligumia subrostrata* releases shell calcium in hypoxic conditions, and it reclaims Ca<sup>2+</sup> as calcium phosphate concretions in the gill tissue.<sup>36)</sup> *P. margaritifera* seemed to need a prolonged time to reclaim Ca<sup>2+</sup> as concretions of calcium phosphate, and [Ca<sup>2+</sup>] slowly decreased as a result, though there was no result of histology in this study. *P. margaritifera* required at least 24 h for the recovery of hemolymph [Ca<sup>2+</sup>].

According to the pH-[HCO<sub>3</sub><sup>-</sup>] diagram of the hemolymph (Fig. 6), [HCO<sub>3</sub><sup>-</sup>] and Pco<sub>2</sub> increased considerably with the reduction in pH, and the points at AE36h were located above the non-bicarbonate buffer line. The buffer value as a measure for the buffering capability is defined as the change in base or acid form of the buffer system per change in pH.<sup>37,38)</sup> The β<sub>NB</sub> is the buffer value of the non-bicarbonate buffer system (mainly protein residues). If a decrease in hemolymph pH is due solely to a change in Pco<sub>2</sub>, the hemolymph will be simply titrated along the non-bicarbonate buffer line,<sup>39)</sup> and the point of pH value moves on this line. If a decrease in hemolymph pH is due solely to an increase in non-volatile acid, then the hemolymph will be titrated along a constant Pco<sub>2</sub> isopleth.<sup>39)</sup> However, *P. margaritifera* showed hemolymph acidosis and high [HCO<sub>3</sub><sup>-</sup>] during air exposure, and the point of AE36h located above the non-bicarbonate buffer line. β<sub>vivo</sub>, which was calculated from hemolymph pH and [HCO<sub>3</sub><sup>-</sup>] at AE0h and AE36h, was 5.22 slykes, and β<sub>vivo</sub> was 10-fold higher than β<sub>NB</sub> (0.53 slykes). Therefore, air-exposed *P. margaritifera* mobilized [HCO<sub>3</sub><sup>-</sup>] from shell carbonate to the hemolymph, and enhanced the buffering capacity of the non-bicarbonate buffer system. Byrne et al. (1991) reported that the resulting base mobilized (primarily bicarbonate) functions to increase the apparent "nonbicarbonate" buffering capacity almost 17-fold over that of isolated hemolymph of the freshwater clam *Corbicula fluminea* during air exposure for 72 h.<sup>27)</sup> This source of readily available buffering power compensates for the low inherent buffering capacity of native hemolymph.<sup>27)</sup> Therefore, *P. margaritifera* enhanced the low buffering capacity of the hemolymph using mobilized base (bicarbonate) from the shell, and provided a partial

metabolic compensation for the acidosis during air exposure in this study.

## References

- 1) Hayami I Pterioidea T: Mytiloidea. In: Okutani T (ed) Marine Mollusks in Japan. Tokai University Press, Tokyo, 879-883 (2000)
- 2) Wada KT, Temkin Ilya: Taxonomy and Phylogeny. In: Southgate PC, Lucas JS (eds) The Pearl Oyster. Elsevier, United Kingdom, 37-75 (2008)
- 3) Yamamoto K, Handa T, Minato Y, Odawara K, Sone K: Effect of low salinity on ventilation volumes in the black-lip pearl oyster *Pinctada margaritifera* and the pearl oyster *Pinctada fucata martensii*. *Aquaculture Sci*, **60**, 277-280 (2012)
- 4) Yamamoto K, Handa T, Araki A, Minato Y, Odawara K and Sone K: Effect of trichlorfon on ventilation in the Black-lip pearl oyster *Pinctada margaritifera*. *J Nat Fish Univ*, **61**, 11-14 (2012)
- 5) Yamamoto K, Handa T, Minato Y, Odawara K, Sone K: Effects of water temperature rise on ventilation in the black-lip pearl oyster *Pinctada margaritifera*. *J Nat Fish Univ*, **61**, 15-18 (2012)
- 6) Yamamoto K, Handa T, Minato Y, Odawara K, Sone K: Effect of hypoxia on oxygen uptake in the black-lip pearl oyster *Pinctada margaritifera*. *J Nat Fish Univ*, **61**, 19-22 (2012)
- 7) Yamamoto K, Handa T: Structure of the digestive diverticula of the black-lip pearl oyster *Pinctada margaritifera* (Mollusca: Bivalvia Pterioidea = Pseudolamellibranchia). *J Nat Fish Univ*, **59**, 39-52 (2010)
- 8) Yamamoto K, Handa T: Structure of the ctenidium and labial palp of the black-lip pearl oyster *Pinctada margaritifera* (Pteriidae: Pterioidea = Pseudolamellibranchia: Bivalvia). *J Nat Fish Univ*, **59**, 53-73 (2010)
- 9) Handa T, Yamamoto K: Estimation of CO<sub>2</sub> partial pressure and bicarbonate concentration in the hemolymph of the black-lip pearl oyster *Pinctada margaritifera*. *J Nat Fish Univ*, **63**, 181-188 (2015)
- 10) Taylor J, Strack E: Pearl production. In: Southgate PC, Lucas JS (eds) The Pearl Oyster. Elsevier, United Kingdom, 273-302 (2008)
- 11) Booth CE, McDonald DG, Walsh PJ: Acid-base balance in the sea mussel, *Mytilus edulis*. I. Effects of hypoxia and air-exposure on hemolymph acid-base status. *Mar Bio Lett*, **5**, 347-358 (1984)
- 12) Michelidis B, Ozounis C, Paleras A, Portner HO: Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Mar Ecol Prog Ser*, **293**, 109-118 (2005)
- 13) Handa T, Yamamoto K: The blood acid-base balance in the pearl oyster, *Pinctada fucata martensii*, after the surgery. *J Nat Fish Univ*, **60**, 57-61 (2011)
- 14) Handa T, Yamamoto K: The acid-base balance of the hemolymph in the pearl oyster *Pinctada fucata martensii* under normoxic conditions. *Aquaculture Sci*, **60**, 113-117 (2012)
- 15) Handa T and Yamamoto K: Estimation of CO<sub>2</sub> partial pressure and bicarbonate concentration in the hemolymph of the noble scallop *Mimachlamys nobilis*. *J Nat Fish Univ*, **64**, 188-194 (2016)
- 16) Handa T, Araki A, Yamamoto K: Acid-base balance of the hemolymph in hard-shelled mussel *Mytilus coruscus* in normoxic conditions. *J Nat Fish Univ*, **65**, 39-46 (2017)
- 17) Handa T, Araki A, Yamamoto K: Oxygen and acid-base status in the Pacific oyster *Crassostrea gigas* after cannulation of the adductor muscle. *J Nat Fish Univ*, **66**, 35-40 (2017)
- 18) Handa T, Araki A, Kawana K, Yamamoto K: Acid-base balance of hemolymph in Pacific oyster *Crassostrea gigas* in normoxic conditions. *J Nat Fish Univ*, **66**, 103-110 (2018)
- 19) Handa T, Araki A, Yamamoto K: Oxygen and acid-base status of hemolymph in the densely lamellated oyster *Ostrea denselamellosa* in normoxic conditions. *J Nat Fish Univ*, **66**, 203-208 (2017)
- 20) Boutilier RG, Iwama GK, Heming TA, Randall DJ: The apparent pK of carbonic acid in rainbow trout blood plasma between 5 and 15 °C. *Resp. Physiol*, **61**, 237-254 (1985)



- 21) Yamamoto K, Adachi S, Tamura I, Aramizu T and Koube H: Effects of hypoxia and water temperature on ciliary movement of gills 5 bivalvia, *Mytilus edulis*, *Atrina pectinata*, *Pinctada fucata martensii*, *Chlamys nobilis* and *Crassostrea gigas*. *J Nat Fish Univ*, **44**, 137-142 (1996)
- 22) Yamamoto K, Handa T: Effect of hypoxia on oxygen uptake in the Pacific oyster *Crassostrea gigas*. *Aquaculture Sci*, **59**, 199-202 (2011)
- 23) Yamamoto K, Handa T, Nakamura M, Kitukawa K, Kita Y, Takimoto S, Nishikawa S: Effects of ozone-produced oxidants on respiration of the pearl oyster, *Pinctada fucata martensii*. *Aquaculture Sci*, **47**, 241-248 (1999)
- 24) Davenport HW: Fundamental equation. *In*: The ABC of acid-base chemistry 6th edition. University of Chicago Press, Chicago, 39-41 (1974)
- 25) Jokumsen A and Fyhn HJ: The influence of aerial exposure upon respiratory and osmotic properties of haemolymph from two intertidal mussels, *Mytilus edulis* L. and *Modiolus modiolus* L. *J Exp Mar Biol Ecol*, **61**, 189-203 (1982)
- 26) Byrne RA, Shipman BN, Smatresk NJ, Dietz and McMahon RF: Acid-base balance during emergence in the freshwater bivalve *Corbicula fluminea*. *Physiol Zool*, **64**, 748-766 (1991)
- 27) Byrne RA and McMahon BR: Acid-base and ionic regulation, during and following emersion, in the freshwater bivalve, *Anodonta grandis simponiana* (Bivalvia: Unionidae). *Biol Bull*, **181**, 289-297 (1991)
- 28) Michaelidis B, Haas D, Grieshaber MK: Extracellular and intracellular acid-base status with regard to the energy metabolism in the oyster *Crassostrea gigas* during exposure to air. *Physiol Biochem Zool*, **78**, 373-383 (2005)
- 29) Handa T, Araki A, Yamamoto K: Effect of air exposure on acid-base balance of hemolymph in hard-shelled mussel *Mytilus coruscus*. *J Nat Fish Univ*, **68**, 65-70 (2020)
- 30) Zwaan AD and Wijsman TCM: Anaerobic metabolism in bivalvia (Mollusca), Characteristics of anaerobic metabolism. *Comp Biochem Physiol*, **54B**, 313-324 (1976).
- 31) Kluytmans, JH, Bont AMT de, Janus J and Wijsman TCM: Time dependent changes and tissue specificities in the accumulation of anaerobic fermentation products in the sea mussel *Mytilus edulis* L. *Comp Biochem Physiol*, **58B**, 81-87 (1977)
- 32) Livingstone DR and Bayne BL: Responses of *Mytilus edulis* L. to low oxygen tension: Anaerobic metabolism of the posterior adductor muscle and mantle tissues. *J Comp Physiol*, **114**, 143-155 (1977)
- 33) Livingstone DR: Anaerobic metabolism in the posterior adductor muscle of the common mussel *Mytilus edulis* L. in response to altered oxygen tension and temperature. *Physiol Zool*, **51**, 131-139 (1978)
- 34) Zwaan A de, Bont AMT de, Zurbug W, Bayne BL and Livingstone DR: On the role of strombine formation in the energy metabolism of adductor muscle of a sessile bivalve. *J Comp Physiol*, **149**, 557-563 (1983)
- 35) Crenshaw MA and Neff JM: Decalcification at the mantle-shell interface in molluscus. *Am Zool*, **9**, 881-885 (1969)
- 36) Silverman H, Steffens WL, Dietz TH: Calcium concretions in the gills of a freshwater mussel serve as calcium reservoir during periods of hypoxia. *J Exp Zool*, **227**, 177-189 (1983)
- 37) Van Slyke DD: On the measurement of buffer values and on the relationship of buffer value to the dissociation constant of the buffer and the concentration and reaction of the buffer solution. *J Biol Chem*, **52**, 525-570 (1922)
- 38) Heisler N: Acid-base regulation. *In*: Evans David H (ed) The physiology of fishes. CRC Press Inc., Florida, 343-378 (1993)
- 39) Wood CM, McMahon BR, McDonald D: An analysis of changes in blood pH following exhausting activity in the starry flounder, *Platichthys stellatus*. *J exp Biol*, **69**, 173-185 (1977)