

Temporal changes in the nutrient status of Matsushima Bay after a wastewater plant was destroyed by a tsunami on 11 March 2011

メタデータ	言語: English
	出版者:
	公開日: 2022-09-28
	キーワード (Ja):
	キーワード (En): Tsunami effect; Wastewater plant;
	Coastal environment; Time trend; Nutrients
	作成者: 奥村, 裕, マスダ, ヨシオ, スズキ, ノリアキ, 筧,
	茂穂, ハラ, モトユキ
	メールアドレス:
	所属: 水産研究・教育機構, 宮城県水産技術総合センター,
	宮城県, 水産研究・教育機構, 東北大学
URL	https://fra.repo.nii.ac.jp/records/72

1	Temporal changes in the nutrient status of Matsushima Bay after a wastewater plant was
2	destroyed by a tsunami on 11 March 2011
3	
4	
5	Yutaka Okumura ^{1*} , Yoshio Masuda ² , Noriaki Suzuki ³ , Shigeho Kakehi ¹ , and Motoyuki
6	Hara ⁴
7	
8	1* Fisheries Resources Institute, National Research and Development Agency, Japan
9	Fisheries Research and Education Agency, 3-27-5, Shinhama, Shiogama, Miyagi 985-
10	0001, Japan
11	2 Miyagi Prefecture Fisheries Technology Institute, Ishinomaki, Japan, 97-6,
12	Sodenohama, Watanoha, Ishinomaki, Miyagi 981-3625, Japan
13	3 Miyagi Prefectural Government, Sendai Regional Promotion Office, Fisheries and
14	Fisheries Port Department, Japan, 1-9-1, Shinhama, Shiogama, Miyagi 985-0001, Japan
15	4 Graduate School of Agricultural Science, Tohoku University, 468-1, Aramaki, Aoba,
16	Aoba, Sendai, Miyagi 980-0845, Japan
17	
18	*Corresponding author: okumura@affrc.go.jp

19 Abstract

We investigated how the nutrient status of Matsushima Bay was affected when a 20wastewater plant was destroyed by a tsunami in March 2011. The nutrient concentrations 21in the seawater and the treated water from the wastewater plant increased just after the 22tsunami, but decreased again to pre-tsunami levels after 2013. The amount of untreated 23water that passed through the wastewater plant decreased just after the tsunami. It was 24estimated that approximately $40.7 \times 10^3 \text{m}^3/\text{d}$ of the treated water from the wastewater 2526plant was discharged to Matsushima Bay; therefore, the quantity of effluent from the wastewater plant was less than one percent of the water inflow from Takagi River and 27Sendai Bay (which are outside Matsushima Bay) to Matsushima Bay. The nutrient 2829concentrations of seawater in Sendai Bay were lower than those in Matsushima Bay. The 30 results suggest that nutrient concentrations in Matsushima Bay after the tsunami did not increase because any untreated or poorly treated effluent was easily diluted by the river 31flow and the inflow of seawater. Many people were concerned about eutrophication, 32therefore, because of the decreased functioning of the wastewater plant in Matsushima 33 Bay. Marked eutrophication in the bay was not observed after August 2011. 34

35

~ ~	T7 1			. 1 .	1	. • .
36	Keywords.	tsunami ettect	wastewater nlant	coastal environment	time frend	nutrients
00	ixcy worus.	isunann encer,	wastewater plant	, coastal chvironnicht.	, unite trenta	, numentos

- 38
- 39

40 Introduction

The 2011 tsunami caused by the Great East Japan Earthquake (Goto et al. 2012), which 41 was the most powerful earthquake ever to hit Japan and had a recurrence interval of 42between 800 and 1,100 years, caused widespread destruction in coastal areas around 43Tohoku. Many researchers have studied the effects of this tsunami on coastal 44 environments and marine organisms in the coastal area around Tohoku, Japan (Kogure et 45al. 2017, Miyazawa et al. 2017, Oozeki et al. 2017, Ito et al. 2018). Some researchers 46 47have also studied how the tsunami affected Matsushima Bay, where there are oyster and seaweed farms and famous tourist spots. Although the tsunami measured in Matsushima 48Bay, which is a closed bay, was lower than in other more exposed coastal areas, it still 49disrupted the sediment on the sea floor through erosion (Okumura and Masuda 2019) and 5051redeposition (Irizuki et al. 2019), which considerably changed the seafloor environment in the bay (Oota et al. 2017), and caused decreases in the mollusc diversity (Sato and 52Chiba 2016) and eelgrass beds (Sakamaki et al. 2016). The earthquake also caused land 53subsidence (Imakiire and Koarai 2012), with the result that the intensity of light reaching 5455the seafloor has decreased, and the eelgrass beds have struggled to recover in some places (Sakamaki et al. 2016). 56

The tsunami destroyed many facilities on land, including wastewater plants. In the 57aftermath of the tsunami, the function of more than 20 wastewater plants in the Tohoku 5859area either temporarily stopped or had decreased function (Matsuhashi et al. 2014, Satoh 2017). Of these wastewater plants, one near Matsushima Bay suffered severe damage 60 (Miyagi Prefectural Government 2011). Other researchers have reported that microbial 61communities in a river that received discharges of sewage effluent changed temporarily 62 63 in the period immediately after the earthquake (Wells et al. 2013). Other researchers reported that the concentrations and composition of nutrients in coastal areas changed 64 after the 2011 tsunami (Fukuda et al. 2016). There was concern that the trophic state of 65

66 Matsushima Bay would increase because of discharges of either untreated or poorly treated wastewater to the bay. While we know that the chemical oxygen demand (COD) 67 and most probable number (MPN) did not change considerably in the months after the 68 tsunami (Sakamaki et al. 2016), there is no information about whether the nitrogen (N) 69 70 and phosphorus (P) concentrations in wastewater discharges emitted to Matsushima Bay changed in that period. N and P serve as indicators of different aspects of water than COD 7172and the MPN (which indicates bacteria sources such as *E.coli*), and increases in the N and 73P concentrations might trigger increases in phytoplankton blooms (USEPA 2004). If the N/P ratio changes, the composition of the phytoplankton assemblages might also change 74(Liu et al. 2013, Xiao et al. 2018) as the demand for nutrients varies with the 7576phytoplankton species. Changes in the phytoplankton species might pose risks for 77 aquaculture; for example, plankton that cause shellfish poisoning would have adverse 78effects on shellfish aquaculture operations (Officer and Ryther 1980); any pause or decrease in the function of a wastewater plant might be harmful for shellfish aquacultures. 79 In this study, we investigated the nutrient concentrations in Matsushima Bay before and 80 81 after the tsunami. We compared information in data reports about the emissions of N and P and quantities of treated water discharged from a wastewater plant in the bay. We also 82 explored how changes in the function of the wastewater plant (from normal to low) 83 affected the nutrient concentrations in the Matsushima Bay, by calculating the water 84 85 quantities, and the N and P concentrations, in the river and in Sendai Bay, into which water from Matsushima Bay flows. 86

87

88 Materials and Methods

89 Investigations

We took samples from 16 stations in Matsushima Bay from August 2011 to December
2015 (Fig. 1). The salinity and water temperature were measured by a Conductivity,

92Temperature, and Depth (CTD) profiler (RINKO-Profiler, JFE-Advantech, Hyogo, Japan). Changes in the seabed topography at each station were measured in meters using 93 the depth sensor feature of the CTD profiler, and this provided information about the 94erosive action of the tsunami and the subsidence caused by the earthquake (Online 95 Resource 1). Samples of seawater to be analyzed for their nutrient concentrations were 96 collected from two depths close to the surface and from 50 cm above the seabed. To 97 calculate how much seawater flowed from the outer area into Matsushima Bay, we also 98 99 collected samples from three stations in Sendai Bay, outside of Matsushima Bay, from 100 April to November (Figs. 1 and 2). These samples were collected from two depths close 101 to the surface and at a depth of 10 m, because the bed at the mouth of Matsushima Bay is 102at about 10 m. A water sample was also collected from the Takagi River, which flows into 103 Matsushima Bay, on seven occasions from 2009 to 2013. The surface water and river water samples were collected in a bucket, and seawater from 50 cm above the seabed was 104 collected with a Van Dorn water sampler (RIGOSHA Co. Ltd., Tokyo, Japan). 105

106

107 Nutrients analysis

The samples were stored in a freezer in the laboratory for approximately 1 month before 108109 analysis. The concentrations of NO₃-N+NO₂-N, NO₂-N, NH₄-N, and PO₄-P in seawater and river water were determined using an Auto Analyzer (BL-TEC, Osaka, Japan), as 110 111 outlined in the instrument's manual. After following the instructions in the instrument's manual, we found the analytical standards of NO₃-N, NO₂-N, PO₄-P, and NH₄-N to be 112KNO₃, NaNO₂, KH₂PO₄, and (NH₄)₂SO₄ (Fujifilm Wako Pure Chemical Co., Osaka, 113Japan), respectively. After weighing each standard chemical, we adjusted the 114concentrations with distilled water. The calibration curve of standards and the 115concentrations of samples were calculated by use of the AACE software (version 6.03) 116 supplied with the Auto Analyzer. The detection limit of each item was less than 0.01 117

118 μ mol/L and is shown as zero in our figures.

119

120 Collecting data from reports

Datasets related to the wastewater plant, such as the concentrations of total N (TN), total 121 P (TP), and various forms of dissolved inorganic nitrogen (DIN; NO₃-N, NO₂-N, and 122NH4-N), in untreated and treated water, and the quantities of wastewater discharged, were 123124obtained from the annual reports of the Miyagi Prefectural Government (2010, 2011, 2012, 1252013, and annual reports from 2014 to 2016 https://www.pref.miyagi.jp/site/sab/list1796-5263.html). In this study, we used data from April 2009 until December 2015. We 126calculated the quantity (Q) of river water using the HQ equation from hourly 127measurements of the river height (H) collected at the Shinainuma Station (Fig. 1) in the 128129Takagi River, obtained from the Miyagi Prefectural Government (unpublished). We 130 obtained data collected before the tsunami about, for example, nutrients and salinities from the Miyagi Prefectural Government (https://www.pref.miyagi.jp/site/kankyo/ in 131132Japanese).

133

134 Calculating the seawater inflow from Sendai Bay to Matsushima Bay

The amount of water that flowed into Matsushima Bay from outside of the bay was 135calculated on the basis of the box model concept (Unoki 1998). The salinity of the 136 137seawater flowing into Matsushima Bay generally decreases when it mixes with freshwater 138 discharged from the river and wastewater plant, but increases when seawater from a deeper layer outside of the bay with high salinity flows into the bay. Therefore, the 139140 amounts of inflow to, and outflow from, Matsushima Bay can be calculated from the salinities and quantities of water from each source, such as the volume in Matsushima 141Bay, wastewater, river water, and inflow from Sendai Bay (Fig. 2). To calculate the inflow 142from Sendai Bay, the volume of Matsushima Bay was calculated, as follows. The bottom 143

144depths measured at the 16 stations were converted to the bottom depth for the average tide level by substituting into the observation time to tide table (because the bottom depth 145changes with the time to the tide). The bottom depth at each station after the tsunami, 146from August 2011 to February 2013, was averaged. We then created a bathymetric map 147of isobath lines of 1 m from the average depths of each station using Ocean Data View 148(https://odv.awi.de/). The areas of each 1-m depth in Matsushima Bay were calculated 149from a bathymetric map from image J (Rasband 2018). Then, the volume in Matsushima 150151Bay was roughly estimated by integrating the area of each 1-m depth.

Inflows from the Takagi River to Matsushima Bay were calculated by substituting the
hourly water heights into the HQ equation. The daily and monthly inflows were calculated
from the hourly inflows.

155To understand the quantities of nutrients that were discharged from the wastewater plant, we investigated temporal trends in the quantities of wastewater and nutrient 156concentrations using monthly data published in the online annual reports from 2009 to 1572015 (Miyagi Prefectural 2010-2013, 158government https://www.pref.miyagi.jp/site/sab/list1796-5263.html). The information for TN, TP, 159NO₃-N, NO₂-N, and NH₄-N was aggregated. We had information about TP but not 160161 dissolved inorganic phosphate (DIP; PO₄-P). Researchers have reported that the DIP/TP ratio for each wastewater plant is different (Tanaka and Shimamura 2005, Yamashita et 162163al. 2006, Kamohara et al. 2019). After referring to an earlier study (Yamashita et al. 2006), 164 we assumed that the DIP/TP ratio was 70%. Treated water from a wastewater plant was discharged to both Sendai Bay and Matsushima Bay through the Sunaoshi-Teizan Canal 165166 in front of a wastewater plant (Figure 1). In 1982 its (Sendai Bay / Matsushima Bay) ratio 167was reported to be 6:4 (Miyagi Prefectural government 1982). The inflow from the wastewater plant to Matsushima Bay was calculated as 40% of the total emissions from 168a wastewater plant. 169

Bi-monthly data for the salinity at the surface and at 50 cm above the seabed were averaged. We averaged the salinities measured at the surface and at a depth of 10 m for each sampling time at the three stations outside of Matsushima Bay (Fig. 1).

173

174 Calculation of DIN and DIP flux from each component

The flux of DIN from Takagi River to Matsushima Bay was calculated by multiplying the river water discharge by DIN concentration in that water. The flux of DIN from the wastewater plant to Matsushima Bay was calculated by multiplying the volume of treated water by DIN concentration in that water; this was further multiplied by 40%. This was because the inflow from the wastewater plant to Matsushima Bay was approximately 40% of the total emissions from that water plant, and 60% outflowed to Sendai Bay (Miyagi Prefectural government 1982).

182 DIP was calculated in the same way as DIN.

183

184 Statistical analyses

Statistical analyses were conducted for the highest or lowest month of nutrient 185concentrations. Significant differences in DIN and DIP concentrations were first 186compared by the Friedman test. Next, nutrient concentrations each year were compared 187 by pairwise comparisons using the Wilcoxon signed-rank test. We used the Bonferroni 188 189 method for P value adjustment. All tests were conducted with EZR (ver. 1.54) software (https://www.jichi.ac.jp/saitama-sct/SaitamaHP.files/statmedEN.html), which is based on 190 R software (https://www.r-project.org/). Pairwise comparisons were not carried out on 191 192samples from Sendai Bay because the sample size was small.

193

194 Results

Temporal trends in the nutrient concentrations

196 Matsushima Bay

197 The mean DIN concentrations tended to vary seasonally through the year, and to be high in autumn (Fig. 3a). The annual maximum value of the monthly means during our survey 198 period was the highest in 2012, and decreased afterwards (Online Resource 2). The annual 199200 maximum value of the monthly means ranged from 7.2 to 16.6 µmol/L before 2011, was 20.4 µmol/L in 2012, and ranged from 10.2 to 14.2 µmol/L after 2013. The annual 201202maximum value in each year was significantly different (p value < 0.05) by the Friedman 203test (Online Resource 3). The mean surface maximum in 2012 was statistically significant in 2009, 2011, 2013, and 2014 at the 0.05 probability level (Online Resource 4a). The 204mean maximum of 50 cm above the seabed in 2012 was statistically significant in all 205206years except in 2010, at the 0.05 probability level (Online Resource 4b).

207The mean DIN minimum in each month in both layers was at its peak in 2011 and 208decreased after 2012 (Fig. 3a). The annual minimum value of the monthly means ranged from 1.9 to 2.7 µmol/L before 2010, from 5.6 to 5.9 µmol/L in 2011, and from 0.14 to 2092100.91 µmol/L after 2012 (Online Resource 2). The annual minimum value in each year 211was significantly different (p value < 0.05) by the Friedman test (Online Resource 3). The mean surface minimum in 2011 was statistically significant after 2012 at the 0.05 212probability level (Online Resource 4c). The mean minimum of 50 cm above the seabed 213in 2011 was statistically significant in all years at the 0.05 probability level (Online 214215Resource 4d). DIN concentrations in Matsushima Bay increased after the tsunami, and tended to be high from 2011 to 2012 and to decrease after 2013. 216

DIP concentrations also tended to fluctuate in cycles in the same way as DIN concentrations (Fig. 3b). Mean DIP concentrations tended to be high in autumn (Online Resource 2). Over the period of the study, the mean DIP maximums in each month in both layers were at their peaks in 2012, and decreased after 2013. The annual maximum value of the monthly means ranged from ranged from 0.77 to 0.9 µmol/L before 2011, from 1.38 μ mol/L to 1.40 μ mol/L in 2012, and from 0.97 to 1.27 μ mol/L after 2013. The annual maximum value in each year was significantly different (*p* value < 0.05) by the Friedman test (Online Resource 3). The mean surface maximum in 2012 was statistically significant in 2009, 2010, 2014, and 2015 at the 0.05 probability level (Online Resource 5a). The mean maximum of 50 cm above the seabed in 2012 was statistically significant in all years at the 0.05 probability level (Online Resource 5b).

The mean DIP minimums in each month in both layers were at their peaks in 2011 and 228229decreased after 2012 (Fig. 3b, Online Resource 2). The annual minimum value of the monthly means ranged from 0.23 to 2.3 µmol/L before 2010, from 0.74 to 5.6 µmol/L in 2302011, and from 0.04 to 0.91 µmol/L after 2012. The annual minimum value in each year 231232was significantly different (p value < 0.05) by the Friedman test (Online Resource 3). The 233mean minimums of the surface and 50 cm above the seabed in 2011 were statistically 234significant in all years at the 0.05 probability level (Online Resources 5c and 5d). DIP concentrations in Matsushima Bay increased after the tsunami, tended to be high from 2352011 to 2012, and to decrease after 2013. 236

The N/P ratio calculated from the average DIN and DIP in each year was 8.7 in 2009,

238 12.4 in 2010, 8.9 in 2011, 11.3 in 2012, 8.3 in 2013, 5.6 in 2014, and 9.5 in 2015.

239

240 Sendai Bay

The DIN concentrations in Sendai Bay also fluctuated throughout the 7 years (Fig. 4a). In contrast to Matsushima Bay, the concentration levels in Sendai Bay varied by year and the month of mean maximum concentrations ranged from April to November; the month of mean minimum concentrations ranged from May to September (Online Resource 6). Time courses of monthly mean maximums were similar to those of Matsushima Bay.

246 The mean DIN maximums in each month in both layers were at their peaks in 2012 and

decreased after 2013. The annual maximum value of the monthly means ranged from 1.19

248 to 3.74 μmol/L before 2012, from 4.88 to 5 μmol/L in 2012, and from 0.55 to 2.95 μmol/L

after 2013. The annual maximum value at a depth of 10 m in each year was significantly different (p value < 0.05) by the Friedman test (Online Resource 3), but that of the surface was not significantly different (p value > 0.05).

The mean DIN minimums in each month in both layers were at their peaks in 2010, and tended to decrease gradually over time. The annual minimum value of the monthly means ranged from 0.62 to 0.64 μ mol/L in 2009, from 1.05 to 0.88 μ mol/L in 2010, and from 0 to 0.55 μ mol/L after 2011. The annual minimum value in each year was significantly different (*p* value < 0.05) by the Friedman test (Online Resource 3).

The monthly mean DIP maximums and minimums resembled those of DIN (Fig. 4). The mean DIP maximums in each month in both layers were at their peaks in 2012, and decreased after 2013 (Online Resource 6). The annual maximum value of the monthly means ranged from 0.13 to 0.42 μ mol/L before 2012, from 0.45 to 0.49 μ mol/L in 2012, and from 0.02 to 0.42 μ mol/L after 2013. The annual maximum value at the two depths were significantly different (*p* value < 0.05) by the Friedman test (Online Resource 3).

263The mean DIP minimums in each month in both layers were at their peaks in 2010, and tended to decrease gradually with time (Fig. 4b, Online Resource 6). The annual 264minimum value of the monthly means were 0 µmol/L in 2009, and ranged from 0.02 to 2650.08 µmol/L in 2010, and from 0 to 0.03 µmol/L after 2011. The periods below the 266267detection limits were 3 months in 2013, 5 months in 2014, and 3-4 months in 2015. The annual maximum value of surface was significantly different (p value < 0.05) by the 268Friedman test, but that at a depth of 10 m was not significantly different (Online Resource 2692703).

The N/P ratio calculated from the average DIN and DIP in each year was 11.3 in 2009,

272 12.2 in 2010, 11.1 in 2011, 9.4 in 2012, 8.6 in 2013, 17.4 in 2014, and 4.2 in 2015.

273

274 Temporal trends in N and P in treated water from the wastewater plant

The TN concentrations doubled just after the tsunami, and then gradually decreased with time and returned to the pre-quake concentrations (Fig. 5a). The annual average concentrations were $1,087 \pm 98 \ \mu mol/L$ in 2009 and $1,062 \pm 62 \ \mu mol/L$ in 2010; they increased to $1,875 \pm 540 \ \mu mol/L$ in 2011 and to $2,019 \pm 273 \ \mu mol/L$ in 2012; decreased to $1,042 \pm 340 \ \mu mol/L$ in 2013 and to $987 \pm 66 \ \mu mol/L$ in 2014; and increased to $1061 \pm 99 \ \mu mol/L$ in 2015.

281The pattern of DIN concentrations changed in a similar way to that of time trends in TN. The annual average DIN concentrations were $1,003 \pm 95 \ \mu mol/L$ in 2009 and 977 ± 52 282 μ mol/L in 2010; they increased to 1,344 \pm 271 μ mol/L in 2011 and to 1,529 \pm 150 μ mol/L 283in 2012; they decreased to $951 \pm 317 \mu mol/L$ in 2013 and to $914 \pm 69 \mu mol/L$ in 2014; 284285and increased to $980 \pm 92 \mu mol/L$ in 2015. The proportions of NH₄-N, NO₂-N, and NO₃-286N varied over time. NH₄-N was approximately 55% of total DIN from April 2009 to February 2011, approximately 100% from April 2011 to December 2012, and 287approximately 55% from January 2013 to February 2015. NH₄-N, therefore, dominated 288289the total DIN in the 2 years after the disaster. NO₃-N was approximately 42% of total DIN from April 2009 to February 2011, approximately 0% from April 2011 to December 2012, 290and approximately 41% from January 2013 to February 2015. NO₃-N was, therefore, 291almost undetected in the 2 years after the tsunami, in contrast to NH₄-N. Levels of NO₂-292293N were always low during these periods, at approximately 3% of total DIN.

The DIN/TN ratio also varied over time. The ratio was approximately 92% from April 2009 to February 2011, 73% from April 2011 until December 2012, and 92% from January 2013 until December 2015. The DIN/TN ratio was low in the 2 years following the tsunami. The average DIN concentrations in the treated water emitted from the wastewater plant were higher than those in seawater of Matsushima Bay (Figs. 3a and 5a). The concentrations in the treated water were 170, 257, and 270 times higher than those in seawater from Matsushima Bay from April 2009 to December 2010, August 2011 to
 December 2012, and from February 2013 to December 2015, respectively.

302 The TP concentration was high just after the tsunami and the changes in TP over time 303 were similar to those of TN (Fig. 5b). The annual average TP concentrations were $52 \pm$ 304 18 μ mol/L in 2009 and 68 \pm 19 μ mol/L in 2010; they increased to 87 \pm 35 μ mol/L in 2011 and to $71 \pm 29 \,\mu$ mol/L in 2012; decreased to $48 \pm 17 \,\mu$ mol/L in 2013; and increased to 65 305 306 \pm 21 µmol/L in 2014 and to 65 \pm 19 µmol/L in 2015. The DIP concentrations were 307 calculated from the TP concentration at 36.4 (52 \times 0.7) μ mol/L in 2009, 47.6 (68 \times 0.7) μmol/L in 2010, 60.9 (87 × 0.7) μmol/L in 2011, 49.7 (71 × 0.7) μmol/L in 2012, 33.6 308 $(48 \times 0.7) \ \mu mol/L$ in 2013, and 45.5 $(65 \times 0.7) \ \mu mol/L$ in 2014 and 2015. The DIP 309 310 concentrations calculated in the treated water were also higher than those in the seawater 311of Matsushima Bay. The concentrations in the treated water were 87, 98, and 95 times 312higher than those in Matsushima Bay seawater from April 2009 to December 2010, from August 2011 to December 2012, and from February 2013 to December 2015, respectively. 313 314 Although TN, TP, and NH₄-N were higher and NO₃-N was lower immediately after the 315earthquake, the levels always fluctuated, and the seasonal changes were unclear.

The N/P ratio calculated from the average DIN and DIP in each year was 27.5 in 2009,

317 20.7 in 2010, 22 in 2011, 30.8 in 2012, 28.3 in 2013, 20 in 2014, and 21.6 in 2015.

318

Raw water inflows to the wastewater plant

The trends in the TN, TP, and NH₄-N concentrations in the raw water inflows to the wastewater plant differed from those in the treated water (Fig. 5c). The annual average concentrations of TN were $3,275 \pm 257 \ \mu mol/L$ in 2009, $3,289 \pm 282 \ \mu mol/L$ in 2010, $2,836 \pm 600 \ \mu mol/L$ in 2011, $3,039 \pm 286 \ \mu mol/L$ in 2012, $3,135 \pm 277 \ \mu mol/L$ in 2013,

- 324 $3,212 \pm 262 \mu mol/L$ in 2014, and $3,087 \pm 289 \mu mol/L$ in 2015. The TN concentrations in
- 325 the raw water were greater than those in the treated water, and were 3.1 times greater from

April 2009 to February 2011, 1.4 times greater from April 2011 to December 2012, and
3.1 times greater from February 2013 to December 2015.

NH₄-N was the main component of the DIN in the raw water, and the concentrations of NO₂-N and NO₃-N were frequently lower than the detection limits of 0.64 and 2.1 μ mol/L, respectively. The trends in the NH₄-N and TN concentrations were similar. The average concentration of NH₄-N was 2,152 μ mol/L from April 2009 to February 2011, 1,765 μ mol/L from April 2011 to December 2012, and then 2,077 μ mol/L. DIN in raw water accounted for approximately 60% of TN through all these periods, but was the main part of TN in the treated water throughout, apart from the period immediately after the tsunami.

336 Nutrient concentrations in the Takagi River

The average concentrations of DIN and DIP in the Takagi River were $19.1 \pm 15.1 \mu mol/L$ and $1.8 \pm 0.6 \mu mol/L$, respectively (Table 1). The nutrient concentrations in the Takagi River were lower than those in the treated water from the wastewater plant and were higher than those in seawater. The average DIN/DIP ratio was 10.5.

341

342 Salinity of water in Matsushima Bay

The salinity tended to vary seasonally (Fig. 6a). The average salinities from May to 343 October ranged from 28.3 to 30.2 PSU at the surface and from 29.1 to 30.9 PSU at 50 cm 344 345above the bottom. The average salinities ranged from 30.9 to 32.1 PSU at the surface, and from 31.6 to 32.3 PSU at 50 cm above the bottom from December to April. The salinities 346 in both layers tended to be lower between May and October than between December and 347 April. The surface salinities generally decreased from June to August and, over this time, 348 349 the difference between the salinity in the surface and 50 cm above the seabed tended to be greater than in other months. 350

351 The salinities of 50 cm above the seabed tended to be slightly higher after the tsunami

352than before the tsunami (Fig. 6a). The salinities from April 2009 to February 2011 ranged from 26.6 to 32.1 PSU at the surface and from 28.3 to 32.3 PSU of 50 cm above the 353 seabed. The salinities at the surface and 50 cm above the seabed from August 2011 to 354December 2012 ranged from 29.3 to 32.7 PSU and from 29.6 to 32.8 PSU, respectively. 355The salinities at the surface and 50 cm above the seabed from February 2013 to December 3562015 were between 21.5 and 32.9 PSU and 27.9 and 33 PSU, respectively. Although the 357 salinity was very low around August 2013, the highest salinity observed between 358February 2013 and December 2015 was higher than the pre-tsunami maximum. 359

360

361 Salinity at the stations in Sendai Bay outside Matsushima Bay

362The salinities at the surface generally decreased from June to September (Fig. 6b). The 363 average salinities at the surface ranged from 26.5 to 33 PSU from April 2009 to November 364 2010, from 26.1 to 33.3 PSU from September 2011 to November 2012, and from 29.6 to 33.3 PSU from April 2012 to November 2015. The average salinities at a depth of 10 m 365 366 ranged from 32.4 to 33.4 PSU from April 2009 to November 2010, from 32 to 33.4 PSU from September 2011 to November 2012, and from 31.8 to 33.4 PSU from April 2012 to 367 368 November 2015. There was little seasonal change in the salinity at a depth of 10 m, which 369 contrasted with the patterns in the surface salinities in Sendai (Fig. 6a) and at both depths 370 measured in Matsushima Bay (Fig. 6b).

371

372 Quantity of water discharged from the wastewater plant

The quantities of treated water discharged from the wastewater plant decreased immediately after the tsunami (Fig. 7a). The average water discharges were 110×10^3 m³/d from April 2009 to February 2011, 56.5 × 10³ m³/d in April 2011, which was approximately half of the average value before the tsunami, 83.6 × 10³ m³/d from April 2011 to December 2012, and then 110×10^3 m³/d, the same as the pre-tsunami level, from January 2013 to February 2015. It took 2 years for the wastewater quantity to recover to
pre-tsunami levels.

When the inflow to Matsushima Bay was assumed to be 40% of the total (Miyagi Prefectural Government 1982), the inflow of treated water to Matsushima Bay averaged 44×10^3 m³/d from April 2009 to February 2011 and from January 2013 to February 2015.

384 Discharge from the Takagi River to Matsushima Bay

385Although there were occasional spikes, the discharge from the river showed a clear seasonal pattern (Fig. 7b). The discharge averaged 1.3×10^6 m³/d from April to September 386 and 0.5×10^6 m³/d for the other periods, from January to March and again from October 387 to December. Although the discharge showed annual variation, the average discharge 388 from April to September was between 1.8 and 3.2 times higher than that from October to 389 the following March. The river water discharge to Matsushima Bay was between 114 390 (5,000/44) to 295 (13,000/44) times greater than the discharge of treated water from the 391392 wastewater plant.

393

Calculating the inflow from Sendai Bay to Matsushima Bay

The inflow from Sendai Bay to Matsushima Bay calculated by the box model varied 395seasonally. The inflow ranged from 3,200 to $115,000 \times 10^3 \text{ m}^3/\text{d}$, and the average was 396 $17,600 \times 10^3 \text{ m}^3/\text{d}$ (Fig. 7c). The average water quantity calculated was 13.5 times higher 397 398 than that from the Takagi River from April to September and was equivalent to approximately 10% of the capacity of Matsushima Bay. The inflow from Sendai Bay to 399 Matsushima Bay tended to increase after the tsunami, although the values varied in 400 different periods. The average inflow was $10,925 \times 10^3 \text{ m}^3/\text{day} \pm 7,200 \times 10^3 \text{ from } 2009$ 401 to 2010, and $19.493 \times 10^3 \text{ m}^3/\text{day} \pm 21.164 \times 10^3$ after 2011. 402

404 Calculating the flow of DIN and DIP in Matsushima Bay

The inflow of DIN to Matsushima Bay was in the order: wastewater plant > Takagi River > Sendai Bay (Figure 8a). The average inflow of DIN was 0.62 ± 0.11 ton/day from the wastewater plant, 0.2 ± 0.1 from Takagi River, and 0.15 ± 0.28 from Sendai Bay. The inflow from the wastewater plant was $3 \times$ and $4 \times$ higher than that from Takagi River and Sendai Bay, respectively.

The inflow of DIP to Matsushima Bay was in the order: wastewater plant \geq Takagi River Sendai Bay (Figure 8b). The average inflow of DIP was 0.06 ± 0.02 ton/day from the wastewater plant, 0.05 ± 0.03 from Takagi River, and 0.04 ± 0.11 from Sendai Bay. However, smaller differences in DIP were found between the inflow from the wastewater plant, Takagi River, and Sendai Bay than were found for DIN.

415

416 Discussion

417 Nutrient sources in Matsushima Bay

The inflow of DIN into Matsushima Bay, ranked from high to low, was highest from the 418 419 wastewater plant, followed by Takagi River, and the lowest was from Sendai Bay (Fig. 8a). DIN concentrations, ranked from high to low, were highest in the wastewater plant, 420followed by the Takagi River and Matsushima Bay, and were lowest in Sendai Bay (Figs. 4213, 4, and 5; Table 1). As the nutrient concentrations were higher at the wastewater plant 422423and in the Takagi River than in Matsushima Bay, we considered that the river and the wastewater plant were the main sources of DIN to Matsushima Bay. We also thought that 424the DIN in Matsushima Bay was mainly diluted by seawater from Sendai Bay, because 425the DIN concentrations in Sendai Bay were lower than those in Matsushima Bay. The 426427results suggest that treated water from the wastewater plant, river water, and seawater were mixed in Matsushima Bay, and then seawater from near to the surface in the bay 428flowed to the outer area of Sendai Bay (Fig. 2). We already know that the nutrient 429

concentrations are influenced by the concentrations outside of the bay through mixing by
estuarine circulation (Yamamoto et al. 2000). If the concentrations outside the bay are
lower than those in the inner areas, such as Matsushima Bay, the concentrations in the
bay are diluted.

The quantities of water flowing into Matsushima Bay were ranked in order, from high 434to low, with most water flowing in from Sendai Bay, followed by the Takagi River, with 435436 the lowest inflow from the wastewater plant (Fig. 7). These results suggest that the 437nutrients and surface salinity in the Matsushima Bay mainly reflect the complex fluctuations and seasonal changes in the seawater inflows from Sendai Bay and the spikes 438in river water, driven by the weather. The low salinities in the surface seawater during 439440 summer reflect the increases in the river water, and thermoclines that develop in the water 441 column.

Various researchers have reported that the seafloor changed because of the tsunamis 442(Murakami et al. 2013, Nishi et al. 2013, Kamiyama et al. 2014, Naiki et al. 2015, Seike 443 et al. 2017, Matsuoka et al. 2018, Okumura et al. 2020), and that the earthquake caused 444 445land subsidence (Imakiire and Koarai 2012, Matsumoto et al. 2018) along the Tohoku Coast. We thought that the inflows from Sendai Bay increased after the earthquake as the 446 bay mouth became deeper and water exchange was easier because of the erosion of the 447bed of Matsushima Bay close to the bay mouth by the tsunami (Okumura and Masuda 448 449 2019) and land subsidence by the earthquake (Sakamaki et al. 2016).

The inflow of DIP into Matsushima Bay tended to be similar to that of DIN, but the influence of DIP from the wastewater plant water was not as high as that of DIN (Fig. 8b). Because the N/P ratio in the treated water was lower than that of seawater and river water, the levels of DIP from the wastewater plant were lower than those of DIN (Figs 3, 4, 5a, and Table 1). We considered, therefore, that the contribution of DIP from the wastewater plant was low. 456

How the damage to the wastewater plant by the tsunami affected the eutrophication in Matsushima Bay

There was concern that the trophic status in the bay would deteriorate because of the 459damage to the wastewater plant. Although the TN and TP concentrations in the treated 460 water from the wastewater plant increased (Fig. 5a, and b), the average DIN and DIP 461 462concentrations in the bay only increased slightly (Fig. 3). We considered that (1) the 463 quantities of wastewater decreased because the houses near the coast were destroyed by 464 the tsunami; (2) the inflows of treated water to Matsushima Bay were limited to 465approximately 40% in all water and the remaining 60% was outflow to Sendai Bay (which 466 is not directly connected to Matsushima Bay) via the canal (Miyagi Prefectural 467 Government 1982, Fig. 7a); (3) the inflow from Sendai Bay was much greater than the 468 inputs of treated water from the wastewater plant or from the river, and the high nutrient concentrations in the treated water were diluted by the seawater from Sendai Bay. While 469 470 the nutrients in the treated water increased after the tsunami, we consider that, because of 471the combination of these factors, the concentrations in Matsushima Bay increased only slightly. 472

The nutrient concentrations in treatment water from the wastewater plant doubled just 473after the tsunami, and then gradually decreased with time and returned to the pre-quake 474475concentrations in 2013 (Fig. 5a and b). With regard to the functioning of sewage treatment 476 plants, the destroyed wastewater plant was restored progressively between March 2011 and April 2013 (Miyagi Prefectural government 2013b). Sedimentation and sterilization 477of wastewater was started in a temporary sedimentation basin from 21 March 2011, about 47847910 days after the tsunami. Simple aeration processing was started by use of a temporary fan machine from 28 June 2011, about 3 months after the tsunami. Microorganism 480 processing was partially started from 4 January 2012. All facilities were restored by 29 481

482 March 2013. Nutrient concentrations decreased with the restoration of the wastewater483 plant.

The TN/DIN and NH₄-N/DIN ratios in the treated water increased just after the tsunami 484 (Fig. 5a). The high NH₄-N/DIN ratio resembled the DIN composition of raw water (Fig. 4855c). These changes may reflect changes in the nitrifying bacteria that convert ammonia 486 to nitrate by nitrification in the wastewater plant (USEPA 2004). Bacteria can convert 487 from nitrate to nitrogen gas by denitrification in a wastewater plant (USEPA 2004). 488 489 Therefore, we considered that the TN/DIN and NH₄-N/DIN ratios changed because of a decrease in the treated capacity of the wastewater plant caused by the decline in the 490 491nitrifying bacteria and other bacteria.

492

493 The influence of the tsunami on the bottom environment

494 The seabed near the mouth of Matsushima Bay was scoured by the tsunami (Irizuki et al. 2019, Okumura and Masuda 2019). After the tsunami, there was less silt in the bottom 495496 sediments and the particles were larger than before, so there were fewer chemicals and 497less oxygen in sediments (Oota et al. 2017). Researchers have reported changes in various sediment properties in Tohoku area, such as the mud contents, COD, acid volatile sulfides 498(Naiki et al. 2015), and organic matter (Nishi et al. 2013), after earthquakes. Sediments 499in the Matsushima Bay environment also improved as a result of outward flow to bottom 500501sediments further offshore through the erosion and resuspension of sediments during the 502tsunami.

When oysters are farmed continuously over a long period, the seabed environment deteriorates because of fecal deposits and other materials (Kimura 1999, Mori 1999, Kawaguchi et al. 2004, Yamamoto et al. 2009). Because of this deterioration, the seabed at the oyster farm was reworked regularly before the earthquake. According to a fisherman, this was no longer necessary after the earthquake. Although N and P are known to leach from the seabed (Kamiyama et al. 1997, Yamamoto et al. 1998), it is possible that the amount of N and P released from the seabed also decreased. These results infer that there were limited increases in the N and P concentrations because of the changes or improvements in the seabed environment.

512

513 Prior to this study, various researchers had reported how the 2011 tsunami influenced 514 organisms and environments, but few had studied how the marine environment changed 515 when the wastewater plants were destroyed. There was great concern that the trophic 516 status would deteriorate rapidly; however, analysis of the nutrient data showed that the 517 nutrients in Matsushima Bay did not increase sharply.

518The TN and DIN concentrations and the TN/DIN and NH₄-N/DIN ratios in the treated 519water changed just after the tsunami because of a decline in the function that caused a 520decrease in the nitrifying bacteria. However, the inflows of treated water to Matsushima Bay were very small compared to those of river water and seawater from outside of the 521bay. Inflows of water with low nutrient concentrations from Sendai Bay increased after 522523the tsunami because of increases in the water exchange, which helps explain why the nutrient concentrations did not increase sharply immediately after the tsunami. Before 524this study, we predicted that eutrophication would progress when a wastewater plant was 525destroyed because of the limited water exchange in closed bays, such as Matsushima Bay. 526527However, we found that eutrophication was prevented after the tsunami by increases in the inflows to Matsushima Bay from outside. About 20 wastewater plants were destroyed 528by the tsunami. Each wastewater treatment plant has different sources and also different 529530capacities for treating water. Bays along the coastline, to which treated wastewater from 531wastewater plants is discharged, may be closed, semi-closed, or open. This study highlights the need for continuous monitoring of coastal environments close to 532wastewater plants. 533

534

535 Acknowledgments

536	This work	was mainly	v supported	by a	grant from	an earthqu	ake damage	recovery	fund
				~		1	0	J	

- 537 from the Fisheries Agency and the Reconstruction Agency, Japan. This was a
- 538 collaborative study between the Tohoku National Fisheries Research Institute and
- 539 Tohoku University. We would like to thank the employees at the wastewater plant for
- 540 their contribution. We thank Edanz Group (https://jp.edanz.com/ac) for editing the
- 541 English text of a draft of this manuscript.

542 References

543	Fukuda H, Katayama R, Yang Y, Takasu H, Nishibe Y, Tsuda A, Nagata T (2016)
544	Nutrient status of Otsuchi Bay (northeastern Japan) following the 2011 off the
545	Pacific coast of Tohoku Earthquake. J Ocean 72:39-52
546	Goto K, Chagué-Goff C, James G, Bruce J (2012) The future of tsunami research
547	following the 2011 Tohoku-oki event. Sediment Geol 282:1-13
548	Imakiire T, Koarai M (2012) Wide-area land subsidence caused by "the 2011 off the
549	Pacific Coast of Tohoku Earthquake." Soils Found 52:842-855
550	Irizuki T, Fujiwara O, Yoshioka K, Suzuki A, Tanaka Y, Nagao M, Kawagata S,
551	Kawano S, Nishimura O (2019) Geochemical and micropaleontological impacts
552	caused by the 2011 Tohoku-oki tsunami in Matsushima Bay, northeastern Japan.
553	Mar Geol 407:261–274
554	Ito K, Fujikura K, Hara M, Aoki M, Ishikawa Y, Sonoda A, Kijima A (2018) Research
555	on effects of the 2011 off the Pacific coast of Tohoku Earthquake and the
556	following tsunami on marine ecosystems to provide scienti ^c information for
557	reconstruction of ^sheries: Comprehensive research by the members of the Tohoku
558	Ecosystem- Assoc. Nippon Suisan Gakkaishi 84:883-884 (in Japanese)
559	Kamiyama T, Naiki K, Watanabe S (2014) Changes in bottom environments and
560	macrobenthic community in embayment of Iwate Prefecture after the
561	tsunamicaused by theGreat East Japan Earthquake. Kaiyo Mon 46:79-85 (in
562	Japanese)
563	Kamiyama T, Tamai K, Tsujino M (1997) Effects of sediment conditions and dissolved
564	oxygen concentration of bottom seawater on nutrient regeneration processes from
565	marine sediments. Bull Nansei Natl Fish Res Inst 30:209-218 (in Japanes with
566	English abstract)
567	Kamohara S, Takasu Y, Yuguchi M, Mima N, Amano Y, Ishida T, Miyawaki D, Suzuki

568	T (2019) Influence of phosphorus management at two wide area sewage treatment
569	plants on water quality in the inner section of Mikawa Bay during the winter
570	season from 2017 to 2018. Aichi Pref Fish Exp Stn Res Rep 13:1-13 (in Japanese
571	with English abstract)
572	Kawaguchi O, Yamamoto T, Matsuda O, Hashimoto T, Takayama H (2004) Artificial
573	midlayer seafloor: simple and new devices to reduce organic loads from oyster
574	rafts to the sediment. Nippon Suisan Gakkaishi 70:722-727 (in Jaoanese with
575	English abstract)
576	Kimura T (1999) The Change of Dissolved and the Decline Oxygen in Oyster Culture
577	Grounds of Growth of Oysters. Aquac Sci 47:119-127 (in Japanese with English
578	abstract)
579	Kogure K, Kawamura T, Nagata T, Hyodo S (2017) Effect off the 2011 Pacific coast of
580	Tohoku Earthquake and the following tsunami on marine ecosystems in Otsuchi
581	Bay and surrounding areas: comprehensive research by the members of the Tohoku
582	Ecosystem-Associated Marine Sciences group (Projet Grand M. Nippon Suisan
583	Gakkaishi 83:642-643 (in Japanese)
584	Liu D, Shen X, Di B, Shi Y, Keesing J, Wang Y (2013) Palaeoecological analysis of
585	phytoplankton regime shifts in response to coastal eutrophication. Mar Ecol Prog
586	Ser 475:1–14
587	Matsuhashi M, Tsushima I, Fukatani W, Yokota T (2014) Damage to sewage systems
588	caused by the Great East Japan Earthquake, and governmental policy. Soils Found
589	54:902–909
590	Matsumoto A, Narita K, Fujita T, Sato T, Matsumoto I, Wada T (2018) Temporal
591	change of eelgrass Zostera marina bed in Matsukawa-ura Lagoon, Fukushima
592	Prefecture. Mer 56:11–20
593	Matsuoka K, Ikeda Y, Kaga S, Kaga Y, Ogata T (2018) Repercussions of the Great East

594Japan Earthquake tsunami on ellipsoidal Alexandrium cysts (Dinophyceae) in Ofunato Bay, Japan. Mar Environ Res 135:123-135 595 Miyagi Prefectural government (1982) Investigation of treated water discharge 596 destination 597 Miyagi Prefectural government (2010) FY2009 Annual report of sewer maintenance in 598Sen-En area. 122 p 599600 Miyagi Prefectural government (2011) FY2010 Annual report of sewer maintenance in 601 Sen-En area. 122 p 602 Miyagi Prefectural government (2012) FY2011 Annual report of sewer maintenance in 603 Sen-En area. 118 p Miyagi Prefectural government (2013a) FY2012 Annual report of sewer maintenance in 604 605 Sen-En area. 122 p Miyagi Prefectural government (2013b) Restoration of the "a revitalized Water" with 606 607 the eyes 608 Miyazawa Y, Tanaka K, Fukuda H, Kida S (2017) A Summary of the Symposium 609 on"Contribution of Coastal Oceanography to Reconstruction following the Great 610 East Japan Earthquake." Bull Coast Ocean Oceanogr 54:95-96 (in Japanes) 611 Mori K (1999) The Present Situation of Self-Pollution of Oyster-Culture Areas and 612 Some Possible Measures to Cope with This Problem. Aquac Sci 47:173-180 (in 613 Japanese with English abstract) 614 Murakami A, Sakai S, Murakami K, Nakamura Y, Okada T, Ttakao T, Shibaki H (2013) Change in Water and Sediment Quality in Ohfunato Bay induced by the 615 616 2011 Great East Japan Tsunami. J Japan Soc Civ Eng Ser B2 (Coastal Eng 617 69:I 496-I 500 (in Japanese with English abstract) Naiki K, Yamada M, Kaga S, Watanabe S, Kamiyama T, Kado R, Ogata T, Nanba N, 618 619 Hayashizaki K-I, Yamada Y, Yamashita T (2015) Spatial and temporal variation of

sediment properties along the coast of the Sanriku District, Iwate Prefecture, due tothe tsunami generated by the Great East Japan Earthquake in 2011. Nippon Suisan

622 Gakkaishi 81:447-455 (in Japanese with English abstract)

- 623 Nishi T, Yagi H, Kamiyama T, Sugimatsu K, Hayashi K, Nakayama A (2013) Change
- 624 of Bottom Environment in Estuaries of Sanriku by Tohoku Earthquake Tsunami. J
- Japan Soc Civ Eng Ser B2 (Coastal Eng 69:I_286-I_290
- 626 Officer CB, Ryther JH (1980) The possible importance of silicon in marine
 627 eutrophication 3:83–91
- 628 Okumura Y, Kaneko K, Ota H, Nagasaka H, Hara M (2020) Analysis of environmental
- and microbiological changes in Onagawa Bay immediately after the tsunami of the
- 630 Great East Japan Earthquake based on sediment cores. Mar Pollut Bull 157:111235
- 631 Okumura Y, Masuda Y (2019) Influence of the tsuinami to the marine environment in
 632 matsushima Bay. AquaNet 22:32-35 (in Japanese)
- 633 Oota H, Suzuki N, Gambe S (2017) Sedimentary Environment before and after the
- Great East Japan Earthquake in Matsushima Bay. Miyagi Pref Rep Fish Sci:35-41
 (in Japanese)
- 636 Oozeki Y, Nakata K, Kishi MJ (2017) Fisheries effects and recovery from the
- 637 earthquake and tsunami of the Great East Japan Earthquake. Fish Oceanogr 26:97–
 638 98
- Rasband W (2018) Image J. Bethesda, Maryland, USA: National Institute of Mental
 Health
- 641 Sakamaki T, Sakurai Y, Nishimura O (2016) Tsunami Impacts on Eelgrass Beds and
- 642 Acute Deterioration of Coastal Water Quality Due to the Damage of Sewage
- 643 Treatment Plant in Matsushima Bay, Japan. Pages 187–199 in V Santiago-Fandiño,
- 644 H Tanaka, M Spiske, eds. Tsunamis and Earthquakes in Coastal Environments.
- 645 Springer International Publishing

646	Sato S, Chiba T (2016) Structural changes in molluscan community over a 15-year
647	period before and after the 2011 great east japan earthquake and subsequent
648	Tsunami around Matsushima Bay, Miyagi prefecture, Northeastern Japan. PLoS
649	One 11:1–10
650	Satoh H (2017) Challenges of Restoring and Rehabilitating Sewer Systems Damaged
651	By the Great East Japan Earthquake and Tsunami. J JSCE 5:279–297
652	Seike K, Shirai K, Murakami-Sugihara N (2017) Using tsunami deposits to determine
653	the maximum depth of benthic burrowing. PLoS One 12:1-11
654	Tanaka T, Shimamura K (2005) Biological Wastewater Treatment Process with
655	Chemical-Phosphorus Recover Reactor 4:101-108 (in Japanese)
656	Unoki S (1998) Relation between the Transport of Gravitational Circulation and the
657	River Discharge in Bays. Oceanogr Japan 7:283-292 (in Japanese with English
658	abstract)
659	USEPA (2004) Primer for Municipal Wastewater Treatment Systems:30
660	Wells NS, Clough TJ, Condron LM, Baisden WT, Harding JS, Dong Y, Lewis GD,
661	Lear G (2013) Biogeochemistry and community ecology in a spring-fed urban
662	river following a major earthquake. Environ Pollut 182:190–200
663	Xiao W, Liu X, Irwin AJ, Laws EA, Wang L, Chen B, Zeng Y, Huang B (2018)
664	Warming and eutrophication combine to restructure diatoms and dinoflagellates.
665	Water Res 128:206–216
666	Yamamoto T, Maeda H, Matsuda O, Hashimoto T (2009) Effects of culture density on
667	the growth and fecal production of the oyster Crassostrea gigas. Nippon Suisan
668	Gakkaishi 75:230-236 (in Japanese with English abstract)
669	Yamamoto T, Matsuda O, Hashimoto T, Imose H, Kitamura T (1998) Estimation of
670	Benthic Fluxes of Dissolved Inorganic Nitrogen and Phosphorus from Sediments
671	of the Seto Inland Sea. Ocean Japan 7:151-158 (in Japanese with English abstract)

672	Yamamoto T, Yoshikawa S, Hashimoto T, Takasugi Y, Matsuda O (2000) Estuarine
673	Circulation Processes in the Northern Hiroshima Bay, Japan.
674	BullCoastOceanogr 37:111–118
675	Yamashita N, Tanaka H, Kiyoshi M, Suzuki Y (2006) Evaluation of effect of sewage
676	effluent on river water quality using AGP and algal growth inhibition tests. JJSCE,
677	SerG 62:191-200 (in Japanese with English abstract)

678

679

681 Figure captions

Fig. 1 Sampling sites (♦: Stations in Matsushima Bay, □: Stations in Sendai Bay, ■: 682 wastewater plant, \triangle : station in Takagi River, \bullet : Shinainuma Station in Takagi River). 683 684 The Takagi River near Shinainuma Station passes under the Naruse River by the siphon. Treated water emitted from a wastewater plant (■) is discharged to the Sunaoshi-Teizan 685Canal, which extends in a north-south orientation. The northern end of the canal is 686 687 connected to Sendai Bay and its southern part is connected to Matsushima Bay. As the 688 inflow ratio of treated water from a wastewater plant to Sendai Bay and Matsushima Bay was reported to be 6:4 in 1982 (Miyagi Prefectural Government 1982), the volume of 689 690 treated water inflow to Matsushima Bay is thought to be approximately 40% to all treated 691 water

692 Fig. 2 Outline of the box model (Unogi 1998). The seawater inflow from Sendai Bay to Matsushima Bay (Q; results are shown in Fig. 7c) can be calculated by using the 693 freshwater quantities (R) from the wastewater treatment plant (results are shown Fig. 7a) 694 695 and Takagi River (results are shown in Fig. 7b), surface salinity in Matsushima Bay (S₁, 696 the result is in Fig. 6a), and salinity at a depth of 10 m in Sendai Bay (S₄, the result is in 697 Fig. 6b). The seawater outflow from Matsushima Bay to Sendai Bay (Q'; results are shown in Fig. 7c) can also be calculated by using the freshwater quantities (R) from the 698 wastewater treatment plant and Takagi River, the salinity of surface in Matsushima Bay 699 700 (S_1) , and the salinity at a depth of 10 m in Sendai Bay (S_4) . S_1 is the average surface 701 salinity at 16 stations in Matsushima Bay (Fig. 1). S₂ is the average salinity of 50 cm above the seabed at 16 stations in Matsushima Bay. S₃ is the average salinity of the surface 702 703 at three stations in Sendai Bay. S₄ is the average salinity at a depth of 10 m at three stations 704in Sendai Bay

Fig. 3 Temporal trends in the DIN and DIP concentrations at two depths in Matsushima
Bay. (a) "Surface" plots are the DIN surface concentrations at 16 stations. "50 cm above

707 the seabed" plots are the DIN concentrations from 50 cm above the seabed at 16 stations. Avg. of 50 cm above the seabed is the average DIN from 50 cm above the seabed at 16 708 stations. Surface Avg. is the average surface DIN at 16 stations. (b) Surface Avg. is the 709 710 average surface DIP at 16 stations. Avg. of 50 cm above the seabed is the average DIP from 50 cm above the seabed at 16 stations. "Surface" plots are the DIP surface 711concentrations at 16 stations. "50 cm above the seabed" plots are the DIP concentrations 712713from 50 cm above the seabed at 16 stations. Arrow indicates March 2011, when the 714tsunami occurred

Fig. 4 Temporal trends in the dissolved inorganic nitrogen (DIN) and dissolved inorganic 715phosphate (DIP) concentrations at two depths in Sendai Bay. (a) "Surface Avg." is the 716average surface DIN at three stations. "Avg. at 10 m depth" is the average DIN from a 717718 depth of 10 m depth at three stations. "Surface" plots are the surface DIN concentrations 719 at three stations. "10m depth" plots are the DIN concentrations from a depth of 10 m depth at three stations. (b) "Surface" plots are the surface DIP concentrations at three 720721stations. "10 m depth" plots are the DIP concentrations from a depth of 10 m depth at 722 three stations. Surface Avg. is the average surface DIP at three stations. "Avg. at 10 m depth" is the average DIP from a depth of 10 m at three stations 723

Fig. 5 Temporal trends in N in treated water discharged from the wastewater plant (a); temporal trends in P in treated water discharged from the wastewater plant (b); and temporal trends in N and P in the inflow to the wastewater plant (c)

Fig. 6 Temporal trends in the salinity concentrations at two depths in Matsushima Bay (a) and Sendai Bay (b). (a) Surface Avg. is the average surface salinity at 16 stations. Avg. of 50 cm above the seabed is the average salinity from 50 cm above the seabed at 16 stations. "Surface" plots are the surface salinities at 16 stations. "50 cm above the seabed" plots are the salinities from 50 cm above the seabed at 16 stations. (b) Surface Avg. is the average surface salinity at three stations. "Avg. at 10 m depth" is the average salinity from a depth of 10 m at three stations. "Surface" plots are the surface salinities at three stations.

"10 m depth" plots are the salinities from a depth of 10 m depth at three stations

Fig. 7 (a) Quantities of water discharged from the wastewater plant. A ratio of 6:4 was 735 736 reported in 1982 for treated water from a wastewater plant that was discharged to Sendai Bay and Matsushima Bay (Miyagi Prefectural Government 1982). Inflow to Matsushima 737 Bay was calculated for 40% of total emissions from the wastewater plant. (b) River water 738 739 discharged from the Takagi River. The water quantities were calculated by substituting river height into the HQ equation in Shinainuma Station. (c) Water quantities from each 740parameter in Matsushima Bay. Outflow is from the upper layer of Matsushima Bay to 741742Sendai Bay. Inflow is the lower layer from Sendai Bay to Matsushima Bay. Takagi River 743is the inflow from Takagi River to Matsushima Bay. Wastewater is the inflow from the 744wastewater plant to Matsushima Bay. Total volume is the seawater volume in Matsushima Bay (unit is $\times 10^3 \text{ m}^3$) 745

Fig. 8 (a) Time courses of total quantities of DIN calculated. "Takagi River" is the inflow of DIN per day (ton/day) from Takagi River to Matsushima Bay. "Wastewater plant" is the inflow of DIN per day in treated water from wastewater plant (ton/day) to Matsushima Bay via the Sunaoshi-Teizan Canal. "Sendai Bay" is the inflow of DIN per day (ton/day) from Sendai Bay to Matsushima Bay. (b) Time courses of total quantities of DIP calculated

- 752
- 753 **Table 1** Nutrient concentrations in the Takagi River
- 754
- Online Resource 1 Water depth in Matsushima Bay (a) before the tsunami; and (b) after
 the tsunami
- 757
- 758